1048-13-151 Marco Fontana\* (fontana@mat.uniroma3.it), Largo S. Leonardo Murialdo, 1, Dipartimento di Matematica, Universita degli Studi Roma Tre, 00146 Rome, Italy, and Gyu Whan Chang, Department of Mathematics, University of Incheon, Incheon, 402-749, South Korea. Star and semistar operations in polynomial extensions. Preliminary report.

The purpose of this talk is to present some of the results contained in two recent papers, joint with G. W. Chang, concerning the problem of extending, in a canonical way, a star or a semistar operation defined on an integral domain D to the polynomial ring D[X]. I will also compare different approaches to this problem considered by several authors with various motivations.

## References

D.D. Anderson, Star operations induced by overrings, Comm. Algebra 16 (1988), 2535-2553.

D.D. Anderson and D.F. Anderson, Examples of star operations on integral domains, Comm. Algebra 18 (1990), 1621-1643.

G.W. Chang and M. Fontana, Uppers to zero and semistar operations in polynomial rings, J. Algebra 318 (2007), 484-493.

G.W. Chang and M. Fontana, Uppers to zero in polynomial rings and Prüfer-like domains, Comm. Algebra 37 (2009), 164-192.

E. Houston, S. Malik, and J. Mott, Characterizations of \*-multiplication domains, Canad. Math. Bull. 27 (1984), 48-52.

A. Mimouni, Note on star operations over polynomial rings, Comm. Algebra 36 (2008), 4249-4256.

G. Picozza, A note on semistar Noetherian domains, Houston J. Math. 33 (2007), 415-432.

P. Sahandi, Semistar-Krull and valuative dimension of integral domains, arXiv:0809.1305, 2008. (Received February

05, 2009)