

GATEAUX DIFFERENTIABLE POINTS WITH SPECIAL REPRESENTATION

SEUNG JAE OH

ABSTRACT. If (x_n) is a bounded sequence in a Banach space, is there an element $x = \sum_{n=1}^{\infty} a_n x_n$ such that $\sum_{n=1}^{\infty} \|a_n x_n\| < \infty$ and the directional derivative of the norm at x , $D(x, x_n)$, exists for every n ? In fact, there are such x 's dense in the closed span of $\{x_n\}$. An application of this fact is made to a proof of Rybakov's theorem on vector measures.

A real-valued function f defined on a linear topological space X is said to be *Gateaux differentiable* at $x \in X$ if, for every $y \in X$,

$$Df(x; y) = \lim_{t \rightarrow 0} \frac{f(x + ty) - f(x)}{t}$$

exists and converges to $x^*(y)$ for a unique $x^* \in X^*$, where X^* denotes the linear space of all continuous linear functionals on X . If $Df(x; y)$ exists for a particular direction $y \in X$, then we call this limit the *Gateaux derivative of f at x in the direction of y* . The notation for the directional derivative of the norm is

$$D(x; y) = \lim_{t \rightarrow 0} \frac{\|x + ty\| - \|x\|}{t}.$$

A real-valued function f on a subset A of a linear space X is said to be *subdifferentiable* at $x \in A$ if there exists $x^* \in X^*$ such that

$$x^*(y - x) \leq f(y) - f(x) \quad \text{for all } y \text{ in } A.$$

We say that x^* is a *subgradient of f at x* . We denote by $\partial f(x)$ the set of all subgradients of f at x and call this set the *subdifferential of f at x* .

THEOREM 1. *Let f be a continuous convex function defined on a Banach space X , and (x_n) a bounded sequence in X ; then there is an element $x = \sum_{n=1}^{\infty} a_n x_n$ such that $\sum_{n=1}^{\infty} \|a_n x_n\| < \infty$ and $Df(x; x_n)$ exists for every n . Further, such x 's are dense in the closed span of x_n 's.*

PROOF. Suppose (x_n) is a bounded sequence in X and f is a continuous convex function on X . Define $T: l_1 \rightarrow X$ by

$$T(a) = \sum_{n=1}^{\infty} a_n x_n \quad \text{where } a = (a_n) \in l_1.$$

Received by the editors March 2, 1984 and, in revised form, June 5, 1984.

1980 *Mathematics Subject Classification.* Primary 46A55; Secondary 46A32, 52A07.

Key words and phrases. Continuous convex function, Gateaux differentiable, countably additive measure.

The norm of T is the surpemum of the norms of x_n 's, so T is a continuous linear operator and $f \circ T$ is a continuous convex function on the separable space l_1 . By Mazur's theorem [4] there is a dense G_δ subset $G \subseteq l_1$ at each point of which $f \circ T$ is Gateaux differentiable. By the continuity of T the set $D = T(G)$ is a dense subset of $T(l_1)$, which is, in turn, dense in the closed span S of $\{x_n\}$. Thus, D is dense in the latter. If x_1^* and x_2^* are elements of the subdifferential $(\partial f)(Ta)$ for some $a \in G$, then

$$x_i^*(x - Ta) \leq f(x) - f(Ta) \quad \text{for all } x \in X, i = 1, 2.$$

In particular,

$$x_i^*(Tb - Ta) \leq f(Tb) - f(Ta) \quad \text{for all } b \in l_1, i = 1, 2,$$

which means $x_1^* \circ T$ and $x_2^* \circ T$ are in $\partial(f \circ T)(a)$. But $f \circ T$ is Gateaux differentiable on G , and, hence, $x_1^* \circ T = x_2^* \circ T$ on l_1 . This implies that $x_1^* = x_2^*$ on $T(l_1)$ and also on S ; that is, f restricted to S is Gateaux differentiable at each point of the dense set D in S since $\partial f(Ta)$ is a singleton set [3, p. 122]. D has the form $\sum_{n=1}^\infty a_n x_n$ with $\sum_{n=1}^\infty \|a_n x_n\| < \infty$, so the proof is complete.

This theorem applies to spaces of measures. Let Σ be a σ -field of subsets of the point set Ω , and let X be a Banach space. The space $ca(\Sigma, X)$ consists of all countably additive X -valued measures on Σ normed by the semivariation. The following is an obvious corollary.

COROLLARY. *If (μ_n) is a bounded sequence in $ca(\Sigma, X)$, then there is an element $\mu = \sum_{n=1}^\infty a_n \mu_n$ such that $\sum_{n=1}^\infty \|a_n \mu_n\| < \infty$ and $D(\mu, \mu_n)$ exists for every n .*

Let $F \in ca(\Sigma, X)$, and let μ be a finite nonnegative real-valued measure on Σ . F is called μ -continuous ($F \ll \mu$) if $\lim_{\mu(E) \rightarrow 0} F(E) = 0$. We now state Rybakov's theorem. ($|\cdot|$ stands for the variation norm.)

THEOREM 2 (RYBAKOV [5]). *If $F \in ca(\Sigma, X)$, then there is an element $x^* \in X^*$ such that $F \ll |x^*F|$.*

In order to prove the theorem we need to use the following Lemma. A proof may be found in [2, pp. 11–13]. We also use the key fact [1] that if $\lambda, \mu \in ca(\Sigma, R)$ and $D(\lambda, \mu)$ exists, then $\mu \ll \lambda$.

LEMMA. *If $F \in ca(\Sigma, X)$, then there exists a nonnegative real-valued countably additive measure μ on Σ such that $F \ll \mu$. Moreover, μ can be chosen so that $\mu = \sum_{n=1}^\infty \beta_n |x_n^*F|$ for some (x_n^*) , with $\|x_n^*\| = 1$ for all n , where the β_n 's can be selected to be nonnegative and satisfy $\sum_{n=1}^\infty \beta_n = 1$.*

PROOF OF THEOREM 2. By using the Lemma let $F \ll \sum_{n=1}^\infty \beta_n |x_n^*F|$, where (x_n^*F) is a bounded sequence in $ca(\Sigma, R)$. Hence, by the Corollary, there is an element μ in $ca(\Sigma, R)$ such that $\mu = \sum_{n=1}^\infty a_n (x_n^*F)$ and $D(\mu, x_n^*F)$ exists for every n . If we set $x^* = \sum_{n=1}^\infty a_n x_n^*$, then $x^*F = \sum_{n=1}^\infty a_n x_n^*F$. Now since $D(x^*F, x_n^*F)$ exists for all n , we have $|x_n^*F| \ll |x^*F|$ for all n [1]. Therefore,

$$F \ll \sum_{n=1}^\infty \beta_n |x_n^*F| \ll |x^*F|.$$

REFERENCES

1. R. G. Bilyeu and P. W. Lewis, *Orthogonality and Hewitt-Yosida theorem in spaces of measures*, Rocky Mountain J. Math. **7** (1977), 629–638.
2. J. Diestel and J. J. Uhl, Jr., *Vector measures*, Amer. Math. Soc., Providence, R. I., 1970.
3. J. R. Giles, *Convex analysis with application in differentiation of convex functions*, Pitman, Boston, Mass., 1982.
4. S. Mazur, *Über konvexe mengen in linearen normierte raumen*, Studia Math. **4** (1933), 70–84.
5. V. Rybakov, *Theorem of Bartle, Dunford, and Schwartz on vector-valued measures*, Mat. Zametki **7** (1970), 247–254.

DEPARTMENT OF MATHEMATICS, HANNAM UNIVERSITY, TAEJEON, CHOONGNAM, KOREA