

SEPARATING VECTORS FOR OPERATORS

D. HAN, D. LARSON, Z. PAN, AND W. WOGEN

(Communicated by Joseph A. Ball)

ABSTRACT. It is an open problem whether every one-dimensional extension of a triangular operator admits a separating vector. We prove that the answer is positive for many triangular Hilbert space operators, and in particular, for strictly triangular operators. This is revealing, because two-dimensional extensions of such operators can fail to have separating vectors.

1. INTRODUCTION

A number of the key motivational examples and pathological counterexamples to open questions in the literature that are contained in the articles [AS1], [AS2], [HLPW], [HLW], [LW1], [LW3], [LW4], [W] have a special structural form which can be described in the following elementary way: They are extensions by algebraic operators (in fact often by one- or two-dimensional operators) of basic types of Hilbert space operators which themselves have especially good (that is, nonpathological) structure. An operator $T \in B(H)$ is called an *extension* of A by C if it has the form

$$T = \begin{pmatrix} A & B \\ 0 & C \end{pmatrix}$$

with respect to an orthogonal decomposition $H = M \oplus N$ for the underlying Hilbert space. The good types that have been studied (so far) are normal operators and triangular operators.

Let \mathcal{A} be an algebra of bounded linear operators on a separable complex Hilbert space H . A vector $x \in H$ is called a *separating vector* for \mathcal{A} if the map $A \rightarrow Ax$, $A \in \mathcal{A}$, is injective. We denote by $Sep(\mathcal{A})$ the set of all separating vectors for \mathcal{A} . For an operator $T \in B(H)$, we use $\mathcal{W}(T)$ to denote the weakly closed unital subalgebra generated by T . An operator T is said to have the *separating vector property*, or simply that T has a *separating vector*, if $\mathcal{W}(T)$ has a separating vector. A vector x in H is called an *algebraic vector* for an operator $T \in B(H)$ if there is a nonzero polynomial p in one variable satisfying $p(T)x = 0$. We use \mathcal{E}_T to denote the set of all algebraic vectors for T .

An operator $T \in B(H)$ is called *triangular* if H has an orthonormal basis $\{e_n : n = 1, 2, \dots\}$ with the property that $Te_n \in span\{e_1, \dots, e_n\}$ for each $n \in \mathbb{N}$.

Received by the editors November 3, 2004 and, in revised form, September 7, 2005.

2000 *Mathematics Subject Classification*. Primary 47A10, 47A65, 47A66, 47B99.

Key words and phrases. Separating vector, extension of operators, triangular operator, integral domain.

The second author was supported in part by NSF grant DMS-0070796.

Equivalently, T is triangular if it has an upper triangular matrix representation for some orthonormal basis indexed by the natural numbers. (See the survey article [H].) It is well known that T is triangular iff the set of algebraic vectors for T is dense in H . An operator $T \in B(H)$ is called *strictly triangular* if it has a strictly upper triangular matrix representation for some orthonormal basis indexed by the natural numbers. In fact, the normal operators that occur in the interesting examples and counterexamples are often diagonal, so are triangular as well, and the triangular operators that occur are often weighted shifts with operator weights, so are strictly triangular, and are sometimes bitriangular operators (i.e. both T and T^* are triangular). These examples in turn motivated interesting research questions for the entire classes of normal operators and triangular operators.

Separating vectors for operators have played a role in several of the papers in the literature, and in particular some of the work of the authors dealing with counterexamples constructed by finite extensions of operators. Separating vectors for operator algebras and linear spaces of operators played an essential role in the work in [L] on algebraic reflexivity. It was a conjecture for awhile that every operator has a separating vector. The work in [W] settled this conjecture negatively, and also answered several longstanding open questions in single operator theory. It was subsequently proven in [GLW] that arbitrary *triangular* operators have separating vectors, and indeed, that $Sep(\mathcal{W}(A))$ is dense in H for every triangular operator.

The example constructed in [W] showing that $\mathcal{W}(T)$ can fail to have a separating vector can be taken to be a two-dimensional extension of a backward shift operator. Therefore the question arises: Does every one-dimensional extension of a triangular operator have the separating vector property? This question was posed in [LW2] and is still open. Let us say that an operator A has *property* (S_n) if $\mathcal{W}(T)$ has a separating vector for all k ($k \leq n$) dimensional extensions T of A . Clearly, property (S_n) implies property (S_k) for all $k \leq n$. In [LW3] and [LW4] it was proven that normal operators and bitriangular operators have property $(S_n), \forall n$.

Suppose that T is strictly triangular. In section 2, we prove Theorem 7, our main result: T has property (S_1) . A key to the proof is that $\mathcal{W}(T)$ is an integral domain. In the last section we study a natural algebra of power series that arises in the proof of Theorem 7. We also consider a density property of the set of separating vectors, and we pose several questions for further study.

2. MAIN RESULTS

In the construction of the counterexamples in [W], $\mathcal{W}(T)$ contains operators in a *corner* of $B(H)$. That is, there are projections P and Q with $PQ = 0$ and so that $PB(H)Q$ meets $\mathcal{W}(T)$. In fact $P\mathcal{W}(T)Q$ can be essentially arbitrarily prescribed, and this is the key to constructing the counterexamples. This motivates the following definition:

If $T \in B(H)$ and if $lat(T)$ denotes the lattice of invariant projections for T , we say that T is *stable* with respect to a projection $P \in lat(T)$ if

$$\mathcal{W}(T) \cap PB(H)P^\perp = (0).$$

This definition of pointwise stability of T at P means precisely that $\mathcal{W}(T)$ contains no (necessarily nonzero) nilpotent operator of index 2 of the form $A = PAP^\perp$. We observe that stability of T at every point P in $lat(T)$ means exactly that $\mathcal{W}(T)$ contains no nonzero nilpotent operator. (Indeed, containment of any nonzero

nilpotent operator implies containment of a nilpotent operator of index 2, and if $A \in \mathcal{W}(T)$ is some nilpotent operator of index 2, then the kernel of A is in $\text{lat}(T)$ and contains the range of A , so letting $P = \text{proj}(\ker(A))$ we see that $A = PAP^\perp$, and thus T is unstable with respect to P . The converse argument is trivial.) Everywhere stability of T is a stronger property than stability at a point of $\text{lat}(T)$. It is the concept of stability at a point that we require in this paper.

Our first result gives a sufficient condition for an extension of an operator by a triangular operator to have the separating vector property and shows that lack of stability is the obstruction to existence of separating vectors.

Suppose that an operator $T \in B(H \oplus K)$ and $P_H \in \text{lat}(T)$, where P_H is the orthogonal projection from $H \oplus K$ onto H . Then T has the form

$$\begin{pmatrix} A & X \\ 0 & B \end{pmatrix}.$$

Recall that for any operator T we use \mathcal{E}_T to denote the set of all algebraic vectors for T .

Proposition 1. *Let T, A, B, X be as above. Suppose that B is a triangular operator and $\mathcal{E}_T = \mathcal{E}_A \oplus 0$. If T is stable with respect to P_H and A has the separating vector property, then so does T .*

Proof. Suppose $B = (b_{ij})$ such that $b_{ij} = 0$ when $i > j$. Let u be a separating vector for $\mathcal{W}(A)$. We show that $u \oplus 0$ is a separating vector for $\mathcal{W}(T)$. Let $S \in \mathcal{W}(T)$ such that $S(u \oplus 0) = 0$. Then S must have the form of

$$\begin{pmatrix} 0 & Y \\ 0 & C \end{pmatrix} = \begin{pmatrix} 0 & y_1 & y_2 & y_3 & \cdot & \cdot & \cdot \\ & c_{11} & c_{12} & \cdot & \cdot & \cdot & \cdot \\ & & c_{22} & \cdot & \cdot & \cdot & \cdot \\ & & & & \cdot & \cdot & \cdot \\ & & & & & \cdot & \cdot \\ & & & & & & \cdot \end{pmatrix}.$$

Here the matrix representations for T and S are with respect to the same decomposition of the underlying space, so one can multiply the operator matrices in the usual fashion.

Let z_i ($i = 1, 2, \dots$) be the column vectors of the matrix $\begin{pmatrix} Y \\ C \end{pmatrix}$. We will show that $z_i \in \mathcal{E}_T$ for all i . If this is done, then, by the assumption that $\mathcal{E}_A \oplus 0 = \mathcal{E}_T$, we will have $c_{ij} = 0$ for all i, j ; hence $C = 0$ and therefore $Y = 0$ since T is stable with respect to P_H . Thus we will obtain $S = 0$ as required.

We have $ST = TS$ because $S \in \mathcal{W}(T)$, so an elementary matrix computation shows that

$$b_{1i}z_1 + \dots + b_{ii}z_i = Tz_i, \quad i = 1, 2, \dots$$

Thus $z_1 \in \mathcal{E}_T$ and so $(T - b_{22}I)z_2 \in \mathcal{E}_T$. Hence $z_2 \in \mathcal{E}_T$. If we have checked that $z_1, \dots, z_{i-1} \in \mathcal{E}_T$, then $(T - b_{ii}I)z_i = b_{1i}z_1 + \dots + b_{i-1,i}z_{i-1} \in \mathcal{E}_T$. So $z_i \in \mathcal{E}_T$, as required. \square

Remark 2. Our main interest in Proposition 1 is the case that A is triangular and not algebraic. For this situation the results of [HLW], and also [HLPW], can be used to construct finite-dimensional extensions to which the proposition can be applied. But note also that if A is any operator such that $\mathcal{E}_A = \{0\}$ (that is, A has no point spectrum), and if $\mathcal{W}(A)$ has a separating vector, then $\mathcal{W}(T)$ has a separating vector for every triangular extension T of A for which $\mathcal{E}_T = \{0\}$.

As usual, a ring \mathcal{A} (in our work \mathcal{A} will be an operator algebra) is an *integral domain* if \mathcal{A} has *no zero divisors* (i.e. if $A, B \in \mathcal{A}$ and $AB = 0$, then either $A = 0$ or $B = 0$.)

Proposition 3. *Let $A \in B(H)$. If $\mathcal{W}(A)$ is an integral domain and has a separating vector, then A has property (S_1) .*

Proof. Let

$$T = \begin{pmatrix} A & b \\ 0 & t \end{pmatrix}$$

be a one-dimensional extension of A with some vector $b \in H$ and some scalar $t \in \mathbb{C}$. Choose $u \in \text{Sep}(\mathcal{W}(A))$. We claim that in fact at most one element in

$$\left\{ \begin{pmatrix} \lambda u \\ 1 \end{pmatrix} : \lambda \in \mathbb{C} \right\}$$

is *not* a separating vector for $\mathcal{W}(T)$. To see this, assume that there exist two different numbers λ_1 and λ_2 such that neither $\lambda_1 u \oplus 1$ nor $\lambda_2 u \oplus 1$ is separating for $\mathcal{W}(T)$. Then there exist operators

$$T_1 = \begin{pmatrix} A_1 & b_1 \\ 0 & t_1 \end{pmatrix}, \quad T_2 = \begin{pmatrix} A_2 & b_2 \\ 0 & t_2 \end{pmatrix}$$

in $\mathcal{W}(T)$ such that $T_i \neq 0$ ($i = 1, 2$), and

$$T_1 \begin{pmatrix} \lambda_1 u \\ 1 \end{pmatrix} = 0, \quad T_2 \begin{pmatrix} \lambda_2 u \\ 1 \end{pmatrix} = 0.$$

Thus

$$T_2 T_1 \begin{pmatrix} \lambda_1 u \\ 1 \end{pmatrix} = 0, \quad T_1 T_2 \begin{pmatrix} \lambda_2 u \\ 1 \end{pmatrix} = 0.$$

Then note that $T_1 T_2 = T_2 T_1$ since $\mathcal{W}(T)$ is abelian. Thus, taking the difference yields

$$T_1 T_2 \begin{pmatrix} (\lambda_1 - \lambda_2)u \\ 0 \end{pmatrix} = 0,$$

which implies that $A_1 A_2 (\lambda_1 - \lambda_2)u = 0$. Since $A_1 A_2 \in \mathcal{W}(A)$ and $u \in \text{Sep}(\mathcal{W}(A))$, it follows that $A_1 A_2 = 0$. Thus either $A_1 = 0$ or $A_2 = 0$ since $\mathcal{W}(T)$ is an integral domain. By assumption we have

$$T_1 \begin{pmatrix} \lambda_1 u \\ 1 \end{pmatrix} = \begin{pmatrix} \lambda_1 A_1 u + b_1 \\ t_1 \end{pmatrix} = 0.$$

So if $A_1 = 0$, then $T_1 = 0$, which contradicts our assumption on T_1 . Similarly, the assumption $A_2 = 0$ gives a contradiction. Thus, except possibly for at most one number λ , $\lambda u \oplus 1 \in \text{Sep}(\mathcal{W}(T))$. \square

Remark 4. Regarding the proof of the above proposition, it is of interest to note that for each $\lambda \in \mathbb{C}$ and each $u \in \text{Sep}(\mathcal{W}(A))$, there is an extension T of A such that $\lambda u \oplus 1 \notin \text{Sep}(\mathcal{W}(T))$. Namely, take

$$T = \begin{pmatrix} A & -\lambda A u \\ 0 & 0 \end{pmatrix}.$$

We give a natural application of Proposition 3, which we will generalize. For a function $\phi \in L^\infty$, we use T_ϕ to denote the Toeplitz operator on H^2 defined by

$$T_\phi f = P\phi f, \quad f \in H^2,$$

where P is the projection from L^2 onto H^2 . Recall that for an operator T , $\{T\}'$ denotes the commutant of T , and $\{T\}''$ denotes the second commutant.

Example 5. *If $\phi \in H^\infty$, then both T_ϕ and T_ϕ^* have property (S_1) .*

Proof. It is well known that

$$\mathcal{W}(T) = \{T_z\}' = \{T_h : h \in H^\infty\}$$

and that the mapping $h \rightarrow T_h$, $h \in H^\infty$, is an (isometric) algebra isomorphism of H^∞ onto $\mathcal{W}(T_z)$. Since H^∞ is an integral domain, so is $\mathcal{W}(T_z)$. Clearly each $f \in H^\infty$ with $f \neq 0$ separates $\mathcal{W}(T_z)$. If $\phi \in H^\infty$, then $\mathcal{W}(T_\phi) \subseteq \mathcal{W}(T_z)$. So we can apply Proposition 3 to conclude that T_ϕ has property (S_1) . Similarly

$$\mathcal{W}(T_\phi^*) = \mathcal{W}(T_\phi)^* \subseteq \mathcal{W}(T_z)^*.$$

Thus $\mathcal{W}(T_\phi^*)$ is an integral domain. Since T_ϕ^* is triangular, $\mathcal{W}(T_\phi^*)$ has separating vectors [GLW]. Thus, by Proposition 3, T_ϕ^* also has property (S_1) . \square

We next show that $\mathcal{W}(T)$ is an integral domain for any nonnilpotent strictly triangular operator.

Lemma 6. *An operator $T \in B(H)$ is strictly triangular if and only if $\bigcup_{n=1}^\infty \ker T^n$ is dense in H .*

Proof. The “only if” direction is clear. The “if” direction follows easily from the observation that the sequence of closed subspaces $\{\ker T^n : 1 \leq n < \infty\}$ is a nest of invariant subspaces for T which has closed union H , for which the restriction of T to each member is nilpotent. \square

Theorem 7. *Every strictly triangular operator has property (S_1) .*

To prove Theorem 7 we need the following results.

Proposition 8. *Let T be a nonnilpotent strictly triangular operator. Then for every operator $A \in \mathcal{W}(T)$, there is a unique formal series*

$$\sum_{k=0}^\infty a_k T^k$$

such that $A|_{\ker T^n} = \sum_{k=0}^n a_k T^k|_{\ker T^n}$ for each n .

Proof. Since T is strictly triangular, by Lemma 6, we have $\bigcup_{n=1}^\infty \ker T^n$ is dense in H . Let $N_1 = \ker T$, and $N_{k+1} = \ker T^{k+1} \ominus \ker T^k$ for all $k \geq 1$. Then $H = \bigoplus_{k=1}^\infty N_k$. If we write $M_n = \bigoplus_{k=1}^n N_k$, then $M_n = \ker T^n$ is an invariant subspace for T . Thus T has the matrix form

$$T = \begin{pmatrix} 0 & T_{12} & & & * \\ 0 & 0 & T_{23} & & \\ 0 & 0 & 0 & T_{34} & \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

such that $T_{k,k+1}$ is one to one for every k because T is not nilpotent. For each n observe that $T|_{M_n}$ is nilpotent of index n .

Let $A \in \mathcal{W}(T)$. Then $A|_{M_n} \in \mathcal{W}(T)|_{M_n} \subseteq \mathcal{W}(T|_{M_n})$. Since $(T|_{M_n})^n = 0$, we have that $A|_{M_n} = p_n(T)|_{M_n}$ for some polynomial p_n of degree $\leq n$. We need to show that for each k , the coefficient $a_k^{(n)}$ of z^k in p_n is independent of $n \geq k$.

Suppose that $p(z) = \sum_{k=0}^m a_k z^k$ is a polynomial of degree $m \geq n$ and $A|_{M_n} = p(T)|_{M_n}$. Let P_k be the projection from H onto N_k . Then

$$P_1 A P_1 = P_1 \sum_{k=0}^m a_k T^k P_1 = a_0 P_1.$$

So a_0 is uniquely determined. Suppose that $j \leq n$ and a_0, a_1, \dots, a_{j-1} have been shown to be independent of p . Then

$$P_1 A P_j = P_1 \sum_{k=0}^j a_k T^k P_j = P_1 \sum_{k=0}^{j-1} a_k T^k P_j + a_j T_{12} T_{23} \dots T_{j,j+1}.$$

Since $T_{1,2} T_{2,3} \dots T_{j,j+1} \neq 0$, we obtain that a_j is uniquely determined. Thus $A = \sum_{k=0}^\infty a_k T^k$, where we interpret this to mean that for any n ,

$$A|_{M_n} = \sum_{k=0}^\infty a_k T^k|_{M_n} = \sum_{k=0}^n a_k T^k|_{M_n}.$$

□

The following is an easy consequence of Proposition 8.

Corollary 9. *Let T be as in Proposition 8.*

- (i) *If $A = \sum_{k=0}^\infty a_k T^k \in \mathcal{W}(T)$, then $T = 0$ if and only if $a_k = 0$ for every $k \geq 0$.*
- (ii) *If $A, B \in \mathcal{W}(T)$ such that $A = \sum_{k=0}^\infty a_k T^k$ and $B = \sum_{k=0}^\infty b_k T^k$, then $AB = \sum_{k=0}^\infty (\sum_{j=0}^k a_j b_{k-j}) T^k$.*

Corollary 10. *If T is a strictly triangular operator which is not nilpotent, then $\mathcal{W}(T)$ is an integral domain.*

Proof. Let $A, B \in \mathcal{W}(T)$ with formal series $A = \sum_{k=0}^\infty a_k T^k$ and $B = \sum_{k=0}^\infty b_k T^k$ and such that $A \neq 0, B \neq 0$. Assume that l (resp. j) is the first nonzero coefficient for A (resp. B). Then, by Corollary 9(ii), AB has a formal series with a nonzero $(l+j)$ -th coefficient. Hence $AB \neq 0$ by Corollary 9(i). Therefore $\mathcal{W}(T)$ has no zero divisors. □

Proof of Theorem 7. *Every strictly triangular operator has property (S_1) .*

Proof. If A is not nilpotent, then it has property (S_1) by Corollary 10 and Proposition 3. If T is nilpotent, then every one-dimensional extension T of A is algebraic. Thus $\mathcal{W}(T)$ has separating vectors. □

Theorem 7 has elementary generalizations to operators T which are strictly lower triangular (the adjoint of a strictly upper triangular operator) as well as to operators which have a strict 2-sided triangular form. By the latter we mean that there is an orthonormal basis for the underlying Hilbert space indexed by the integers, $\{e_n : n \in \mathbb{Z}\}$, such that $T e_n \in [e_k : k < n]$ for all n , where $[\]$ denotes a closed linear span. In [GLW, Corollary 12] it was proven that triangular, lower triangular, and 2-sided triangular operators have the separating vector property.

Proposition 11. *If $T \in B(H)$ is either strictly triangular, strictly lower triangular, or has a strict 2-sided triangular form, then either T is nilpotent or $\mathcal{W}(T)$ has no zero divisors. Hence T has property (S_1) .*

Proof. Suppose that T is *not* nilpotent. If T is strictly triangular, then $\mathcal{W}(T)$ has no zero divisors by Corollary 10. If it is strictly lower triangular, then $\mathcal{W}(T^*)$ is an integral domain, hence so is $\mathcal{W}(T) = (\mathcal{W}(T^*))^*$.

The case remains where T has a strict 2-sided triangular form. Let $\{e_n : n \in \mathbb{Z}\}$ be the corresponding orthonormal basis for H , and for each n let P_n be the orthogonal projection onto $E_n := [e_k : k \leq n]$. Then each compression $P_n T|_{E_n}$ is strictly lower triangular and each compression $P_n^\perp T|_{E_n^\perp}$ is strictly triangular. If both of these compressions were nilpotent, then T would have the form

$$\begin{pmatrix} A & X \\ 0 & B \end{pmatrix},$$

where A and B are nilpotent, and this would imply that T was nilpotent, a contradiction. Thus for each n at least *one* of these two compressions is *not* nilpotent. A similar operator matrix argument shows that if $P_n T|_{E_n}$ is not nilpotent, then $P_m T|_{E_m}$ is not nilpotent for all $m \in \mathbb{Z}$, and if $P_n^\perp T|_{E_n^\perp}$ is not nilpotent, then $P_m^\perp T|_{E_m^\perp}$ is not nilpotent for all $m \in \mathbb{Z}$.

Assume, by way of contradiction, that $\mathcal{W}(T)$ is not an integral domain. Then there exist $A_1, A_2 \in \mathcal{W}(T)$, $A_i \neq 0$, $A_1 A_2 = 0$. We have two cases:

Case I. $P_0 T|_{E_0}$ is not nilpotent. Since $P_n \rightarrow I$ strongly as $n \rightarrow \infty$ there exists $k \in \mathbb{Z}$ such that $P_k A_i|_{E_k} \neq 0$, $i = 1, 2$. We also have $P_k \mathcal{W}(T)|_{E_k} \subseteq \mathcal{W}(P_k T|_{E_k})$. Moreover $P_k T|_{E_k}$ is not nilpotent; hence $\mathcal{W}(P_k T|_{E_k})$ is an integral domain by the first paragraph. However,

$$P_k A_1|_{E_k} \cdot P_k A_2|_{E_k} = P_k A_1 A_2|_{E_k} = 0,$$

which is a contradiction. Hence $\mathcal{W}(T)$ must be an integral domain.

Case II. $P_0^\perp T|_{E_0^\perp}$ is not nilpotent. Since $P_n^\perp \rightarrow I$ strongly as $n \rightarrow -\infty$ there exists $k \in \mathbb{Z}$ such that $P_k^\perp A_i|_{E_k^\perp} \neq 0$, $i = 1, 2$. The rest of the argument is analogous to Case I. □

The above result can be extended to the following more general situation.

Theorem 12. *Let $\{A_i \in B(H_i) : i = 1, 2, \dots\}$ be a sequence of operators such that either A_i is algebraic or $A_i = T_i + c_i I$ for some strictly triangular, strictly lower triangular or strictly 2-sided triangular operator T_i . Then $A := \bigoplus_{i=1}^\infty A_i$ has property (S_1) .*

To prove Theorem 12 we need the following lemma.

Lemma 13. *Let $A_i \in B(H_i)$ and*

$$\hat{A}_i = \begin{pmatrix} A_i & x_i \\ 0 & 0 \end{pmatrix} \in B(H_i \oplus \mathbb{C})$$

such that $0 \oplus 1 \in \overline{\text{Sep}(\mathcal{W}(\hat{A}_i))}$ and $X = (x_1, x_2, \dots) \in \bigoplus_i H_i$. Let

$$\hat{A} = \begin{pmatrix} A & X^t \\ 0 & 0 \end{pmatrix},$$

where $A = \bigoplus_{i=1}^\infty A_i$. Then $\mathcal{W}(\hat{A})$ has a separating vector.

Proof. Since $0 \oplus 1 \in \overline{\text{Sep}(\mathcal{W}(\hat{A}_i))}$, we can choose $u_i \oplus 1 \in \text{Sep}(\mathcal{W}(\hat{A}_i))$ such that $\sum_i \|u_i\|^2 \leq \infty$. We claim that $v = (\bigoplus_i u_i) \oplus 1 \in \text{Sep}(\mathcal{W}(\hat{A}))$. To this purpose, let

$\hat{B} \in \mathcal{W}(\hat{A})$ such that $\hat{B}v = 0$. We can write $\hat{B} = \begin{pmatrix} B & Y \\ 0 & 0 \end{pmatrix}$ with $B = \bigoplus_{i=1}^{\infty} B_i$ and $Y = (y_1, y_2, \dots)^t$. Let

$$\hat{B}_i = \begin{pmatrix} B_i & y_i \\ 0 & 0 \end{pmatrix}.$$

Then $\hat{B}_i \in \mathcal{W}(\hat{A}_i)$ and $\hat{B}_i(u_i \oplus 1) = 0$, which implies that $\hat{B}_i = 0$. Thus $\hat{B} = 0$. \square

Proof of Theorem 12.

Proof. Let $\hat{A} = \begin{pmatrix} A & X^t \\ 0 & \lambda \end{pmatrix}$ be any one-dimensional extension of A . Write $X = (x_1, x_2, \dots)$. By replacing \hat{A} by $\hat{A} - \lambda I$, noting that the new operator has the same form, we can assume that $\lambda = 0$. If A_i is algebraic, then so is \hat{A}_i , where $\hat{A}_i = \begin{pmatrix} A_i & x_i \\ 0 & 0 \end{pmatrix}$. Thus $\text{Sep}(\mathcal{W}(\hat{A}_i))$ is dense in $H_i \oplus \mathbb{C}$. If $A_i = T_i + c_i I$ is not algebraic, then $\mathcal{W}(A_i)$ is an integral domain by Proposition 11. Therefore, by the proof of Proposition 3, $0 \oplus 1 \in \overline{\text{Sep}(\mathcal{W}(\hat{A}_i))}$. So the conclusion follows from Lemma 13. \square

Corollary 14. *Suppose that $A_i \in B(H_i)$ such that either A_i is algebraic or $\mathcal{W}(A_i)$ is an integral domain for all $i = 1, 2, \dots$. If each A_i has property (S_1) , then so does $\bigoplus_{i=1}^{\infty} A_i$.*

Note that $\mathcal{W}(\bigoplus_i A_i)$ is not necessarily an integral domain in general. Thus the above corollary is an extension of Proposition 3.

Proof. If A_i is algebraic, clearly $0 \oplus 1 \in \overline{\text{Sep}(\mathcal{W}(\hat{A}_i))}$. If $\mathcal{W}(A_i)$ is an integral domain, then, from the proof of Proposition 3, $0 \oplus 1 \in \text{Sep}(\mathcal{W}(\hat{A}_i))$. Hence $\bigoplus_i A_i$ has property (S_1) by Lemma 13. \square

Remark 15. Proposition 11 is the best possible general result of its kind for operators with a triangular form modeled on a nest of invariant subspaces. The reason is that strictly triangular operators can fail to have property (S_2) , and the example in [W] which points this out (although of course the terminology is different) is a 2-dimensional extension T of a strictly triangular operator A such that T itself has a strictly triangular form with respect to the associated nest order-isomorphic to $\mathbb{N} + \{1, 2\}$. So the intermediate operator B , which is the corresponding 1-dimensional extension of A , has strictly triangular form with respect to $\mathbb{N} + \{1\}$, yet *cannot* be an integral domain operator because B has a 1-dimensional extension, namely T , which does not have the separating vector property.

3. RELATED RESULTS AND QUESTIONS

We saw in Proposition 8 that if T is strictly triangular, then each element $A \in \mathcal{W}(T)$ is a formal power series

$$A = \sum_{k=0}^{\infty} a_k T^k.$$

We can interpret this equality as saying that for any n ,

$$(*) \quad Ae_n = \sum_{k=0}^{\infty} a_k T^k e_n,$$

where the sum on the right side has only finitely many terms which are nonzero. It is natural to define

$$\mathcal{F}(T) = \{A \in B(H) : A = \sum a_k T^k \text{ formally as in } (*)\}.$$

Versions of Corollaries 9 and 10 hold for $\mathcal{F}(T)$, and one can show the following.

Proposition 16. *Let T be strictly triangular. Then:*

- (i) $\mathcal{F}(T)$ is a weakly closed abelian algebra.
- (ii) $\mathcal{W}(T) \subset \mathcal{F}(T) \subset \{T\}''$.
- (iii) If T is not nilpotent, then $\mathcal{F}(T)$ is an integral domain.

We illustrate the containments in Proposition 16(ii) by the following examples.

Example 17. Suppose that T is a backward unilateral weighted shift with all weights nonzero. It is well known [S] that $\{T\}'$ is an algebra of power series in T and that the Cesaro sums of each such series converge weakly. Hence $\mathcal{W}(T) = \mathcal{F}(T) = \{T\}'' = \{T\}'$.

Example 18. We give an example showing that $\mathcal{F}(T)$ may be much larger than $\mathcal{W}(T)$ and that the formal series of an operator in $\mathcal{F}(T)$ need not converge in any sense. Let $G = \{z : 1 < |z| < 3\}$, and let m be the area measure on G . Define

$$L_a^2(G) = \{f : G \rightarrow \mathbb{C} \mid f \text{ is analytic on } G \text{ and } f \in L^2(m)\}.$$

If $\phi \in H^\infty(G)$, let $S_\phi f = \phi f$, $f \in L_a^2(G)$. Then S_z is a subnormal bilateral weighted shift [S] on $L_a^2(G)$. Fix $\lambda \in G$ and let $T^* = S_z - \lambda I$. Then $f \in \text{ran}(T^{*k})$ if and only if f has a zero of order at least k at λ . Thus $\bigcap_{k=1}^\infty \text{ran}(T^{*k}) = \{0\}$, so $\bigcup_{k=1}^\infty \text{ker} T^k$ is dense. Therefore T is strictly triangular.

Now $\mathcal{W}(T^*) = \{S_\phi : \phi \in H^\infty(\{z : |z| < 3\})\}$, while $\{T^*\}' = \{S_\phi : \phi \in H^\infty(G)\}$. Thus $\mathcal{W}(T^*) \subsetneq \{T^*\}'$, and therefore $\mathcal{W}(T) \subsetneq \{T\}'$. Furthermore if $\text{ker} T^k = M_k$, then M_k is invariant for $\{T\}'$ (that is, M_k is hyperinvariant) for each k , and $T|_{M_k}$ is nilpotent of index k . Thus $T|_{M_k}$ is similar to a nilpotent Jordan block J_k on \mathbb{C}^k . (Here $J_k e_1 = 0$ and $J_k e_n = e_{n-1}$, $2 \leq n \leq k$.) But $\{J_k\}' = \{p(J_k) : p \text{ is a polynomial of degree } \leq k\}$. Thus if $A \in \{T\}'$, then $A|_{M_k} \in \{T|_{M_k}\}'$.

Therefore $A|_{M_k} = p_k(T)|_{M_k}$ for some polynomial p_k . It is easy to see (see the proof of Proposition 8) that the coefficients of p_k are uniquely determined, and that $A = \sum_k a_k T^k$, formally. Thus $\mathcal{W}(T) \subsetneq \mathcal{F}(T) = \{T\}' = \{T\}''$.

Example 19. In this example we outline the construction of a strictly triangular operator S so that $\mathcal{W}(S) = \mathcal{F}(S)$ is properly contained in $\{S\}''$. Suppose that S_1 and S_2 are backward weighted shifts on l^2 with nonzero weights, and let $S = S_1 \oplus S_2$. Then S is an operator weighted shift, and, as in the scalar case, one has that $\mathcal{W}(S)$ is an algebra of power series in S , so that $\mathcal{W}(S) = \mathcal{F}(S)$. Now suppose that the weights have been chosen so that there are no nonzero operators intertwining the two summands of S . It follows that $\{S\}' = \{S_1\}' \oplus \{S_2\}' = \{S\}''$, which properly contains $\mathcal{W}(S)$.

One can construct S_1 and S_2 as follows. We will choose both sequences of weights to be constant on blocks of size $2, 2^2, 2^3, \dots$. For S_1 , let the weights be 1 on all of the odd blocks and $1/2$ on the even blocks. For S_2 , the weights will be 1 on the even blocks and $1/2$ on the odd blocks. A matrix computation shows that if $AS_1 = S_2A$ or if $AS_2 = S_1A$, then $A = 0$.

Note that with T as in Example 18, the operator $S \oplus T$ is strictly triangular and $\mathcal{F}(S \oplus T)$ lies properly between $\mathcal{W}(S \oplus T)$ and $\{S \oplus T\}''$.

We say that a nonempty set E of H is *linearly dense* in H if $E \cap L$ is dense in L for all complex lines L that meet E . (By a *complex line* we mean a one-dimensional complex affine subspace of H .) This is a stronger property than density. It was proven in [GLW] that if \mathcal{A} is a linear subspace of operators with denumerable Hamel basis, then $Sep(\mathcal{A})$ is either empty or linearly dense. Also, if \mathcal{A} is a m.a.s.a, then $Sep(\mathcal{A})$ is linearly dense. We conjecture that linear density is common when $Sep(\mathcal{A})$ is nonempty.

Question 1. If $\mathcal{A} \subseteq B(H)$ is an integral domain and $Sep(\mathcal{A})$ is nonempty, is $Sep(\mathcal{A})$ linearly dense?

The following elementary result shows the above is true when \mathcal{A} is abelian.

Proposition 20. *If \mathcal{A} is an abelian integral domain and $Sep(\mathcal{A})$ is nonempty, then $Sep(\mathcal{A})$ is linearly dense.*

Proof. Let $u \in Sep(\mathcal{A})$ and let $x \in H$ be an arbitrary element. Assume that there exist two different scalars $\lambda_1, \lambda_2 \in \mathbb{C}$ such that $u + \lambda_1 x, u + \lambda_2 x \notin Sep(\mathcal{A})$. Then there exist nonzero operators $A_1, A_2 \in \mathcal{A}$ such that $A_1(u + \lambda_1 x) = 0$ and $A_2(u + \lambda_2 x) = 0$. Note that $A_1 A_2 = A_2 A_1$. We have $(\lambda_1 - \lambda_2) A_1 A_2 u = 0$. Hence $A_1 A_2 = 0$, which implies that either $A_1 = 0$ or $A_2 = 0$. Therefore except for possibly at most one λ , all $u + \lambda x \in Sep(\mathcal{A})$, which implies that $Sep(\mathcal{A})$ is linearly dense. \square

Remark 21. The separating vector index was introduced in [HLW] which generalizes the concept of spectrum cardinality for operators acting on finite-dimensional Hilbert spaces. Let \mathcal{A} be a linear subspace of operators such that $Sep(\mathcal{A})$ is nonempty. Define

$$i(\mathcal{A}; L) = \text{card}\{y \in L : y \notin Sep(\mathcal{A})\}$$

for any complex line L meeting $Sep(\mathcal{A})$, and for any $x \in Sep(\mathcal{A})$ define

$$j(\mathcal{A}; x) = \sup\{i(\mathcal{A}; L) : L \text{ is a complex line containing } x\}.$$

The *separating vector index* of \mathcal{A} is defined by $k(\mathcal{A}) = \sup_x j(\mathcal{A}; x)$. If T acts on a finite-dimensional space, then $k(\mathcal{W}(T))$ is the cardinality of the spectrum of T ([GLW]). Proposition 20 tells us that the separating vector index of \mathcal{A} is at most 1 when \mathcal{A} is an abelian integral domain which has a separating vector.

Question 2. Let \mathcal{A} be a weakly closed subalgebra of $B(H)$. If \mathcal{A} is an integral domain, does \mathcal{A} have a separating vector?

Note that Wogen's example [W] shows that property (S_1) does not imply property (S_2) . We have:

Question 3. If T has property (S_n) ($n \geq 2$), does it have property (S_{n+1}) ?

Question 4. Assume the hypotheses of Proposition 1, and suppose that K has dimension 1. Must T have a separating vector that is not in H ? A counterexample would provide the first known example of an operator T with a separating vector such that $Sep(\mathcal{W}(T))$ is not dense. The question is open even for the case that A is triangular. A counterexample in this setting would provide evidence that there may be a one-dimensional extension of a triangular operator that has no separating vector.

REFERENCES

- [AS1] E. Azoff and H. Shehada, *Algebras generated by mutually orthogonal idempotent operators*, J. Operator Theory, **29** (1993), 249-267. MR1284883 (95h:47063)
- [AS2] E. Azoff and H. Shehada, *Literal embeddings of linear spaces of operators*, Indiana Univ. Math. J., **42** (1993), 571-589. MR1237060 (94i:47068)
- [Co] J. Conway, *A Course in Functional Analysis*, Springer-Verlag, New York Inc., 1985. MR0768926 (86h:46001)
- [GLW] W. Gong, D. Larson and W. Wogen, *Two results on separating vectors*, Indiana University Math. J., **43**(1994), 1159-1165. MR1322615 (96h:47050)
- [HN] D. Hadwin and E. Nordgren, *Subalgebras of reflexive algebras*, J. Operator Theory, **7**(1982), 3-23. MR0650190 (83f:47033)
- [HLPW] D. Han, D. Larson, Z. Pan and W. Wogen, *Extensions of operators*, Indiana University Math. J., **53** (2004), no. 4, 1151-1169. MR2095452
- [H] D. Herrero, *Triangular Operators*, Bull. London Math. Soc. **23**(1991), 513-554. MR1135184 (92k:47001)
- [HLW] D. Herrero, D. Larson and W. Wogen, *Semitriangular operators*, Houston J. Math., **17**(1991), 477-499. MR1147270 (92m:47037)
- [L] D. R. Larson, *Reflexivity, algebraic reflexivity and linear interpolation*, Amer. J. Math., **110**(1988), 283-299. MR0935008 (89d:47096)
- [LW1] D. Larson and W. Wogen, *Reflexivity properties of $T \oplus 0$* , J. Funct. Anal., **92**(1990), 448-467. MR1069253 (91i:47010)
- [LW2] D. Larson and W. Wogen, *Some problems on triangular and semitriangular operators*, Contemporary Mathematics, **120**(1991), 97-100. MR1126279 (92i:47019)
- [LW3] D. Larson and W. Wogen, *Extensions of normal operators*, Integr. Equat. Oper. Th., **20**(1994), 325-334. MR1299891 (96b:47024)
- [LW4] D. Larson and W. Wogen, *Extensions of bitriangular operators*, Integr. Equat. Oper. Th., **25**(1996), 216-223. MR1388681 (97d:47027)
- [S] A. Shields, *Weighted shift operators and analytic function theory*, Topics in Operator Theory, Math. Surveys, Amer. Math. Soc., Providence R. I. (1974), no. 13. MR0361899 (50:14341)
- [W] W. Wogen, *Some counterexamples in nonselfadjoint algebras*, Ann. Math., **126** (1987), 415-427. MR0908152 (89b:47066)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CENTRAL FLORIDA, ORLANDO, FLORIDA 32816
E-mail address: dhan@pegasus.cc.ucf.edu

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843
E-mail address: larson@math.tamu.edu

DEPARTMENT OF MATHEMATICS, SAGINAW VALLEY STATE UNIVERSITY, UNIVERSITY CENTER, MICHIGAN 48710
E-mail address: Pan@svsu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NORTH CAROLINA 27599
E-mail address: wrw@math.unc.edu