

LARGE TIME BEHAVIOR AND GLOBAL EXISTENCE OF SOLUTION TO THE BIPOLAR DEFOCUSING NONLINEAR SCHRÖDINGER-POISSON SYSTEM

BY

CHENGCHUN HAO AND LING HSIAO

Academy of Mathematics and Systems Science, CAS, Beijing, 100080, P. R. China

Abstract. In this paper, we study the large time behavior and the existence of globally defined smooth solutions to the Cauchy problem for the bipolar defocusing nonlinear Schrödinger-Poisson system in the space \mathbb{R}^3 .

1. Introduction. In the present paper, we study the global existence and large time behavior for the bipolar defocusing nonlinear Schrödinger-Poisson (BDNLSP) system

$$i\varepsilon\dot{\psi}_j = -\frac{\varepsilon^2}{2}\Delta\psi_j + (q_jV + h_j(|\psi_j|^2))\psi_j, \quad j = 1, 2, \quad (1.1)$$

$$-\lambda^2\Delta V = |\psi_1|^2 - |\psi_2|^2, \quad (1.2)$$

with the initial data

$$\psi_j(0, \cdot) = \varphi_j, \quad j = 1, 2, \quad (1.3)$$

where the wave function $\psi_j = \psi_j(t, x) : \mathbb{R}^{1+d} \rightarrow \mathbb{C}$, $j = 1, 2$, $\dot{\psi}_j = \partial\psi_j/\partial t$, Δ is the Laplace operator on \mathbb{R}^d , and the electrostatic potential $V = V(t, x)$. The nonlinear self-interacting potential $h_j(s)$ is assumed to be given by

$$h_j(s) = a_j^2 s^{\gamma_j}, \quad \text{for } s \geq 0 \text{ and some } a_j > 0, \quad \frac{2}{d} < \gamma_j < \alpha(d),$$

where $\alpha(d) = \frac{2}{d-2}$ if $d \geq 3$ and $\alpha(d) = \infty$ if $d = 1, 2$. The charges of the particles described by the wave functions ψ_j are defined by $q_1 = 1$, $q_2 = -1$, respectively. ε is the scaled Planck constant and λ is the scaled Debye length.

Received July 31, 2003.

2000 *Mathematics Subject Classification.* Primary 35Q55, 82D10.

Key words and phrases. Bipolar defocusing nonlinear Schrödinger-Poisson system, global existence, large time behavior, pseudo-conformal conservation law, the Galilei-type operator.

L. Hsiao is supported by the MST (Grant No. 1999075107), NSAF (Grant No. 10276036), and the Innovation funds of AMSS, CAS of China.

E-mail address: hcc@mail.amss.ac.cn

E-mail address: hsiao1@mail.amss.ac.cn

We assume that the initial value

$$\varphi_j(x) \in \Sigma(\mathbb{R}^d) := \{u \in H^1(\mathbb{R}^d) : |x|u \in L^2(\mathbb{R}^d)\}, \quad j = 1, 2, \quad (1.4)$$

with the norm

$$\|\psi_j\|_{\Sigma} = \|\psi_j\|_{H^1} + \| |x| \psi_j \|_{L^2}.$$

This system appears in quantum mechanics as well as semi-conductor and plasma physics. A large amount of interesting work has been devoted to the study of the Schrödinger-Poisson systems (see [2], [3], [4], [6], [7] and references therein). In [4], by applying the estimates of a modulated energy functional and the Wigner measure method, Jüngel and Wang discussed the combined semi-classical and quasineutral limit of the (BDNLSP) system with the initial data (1.3) in the whole space where $a_1 = a_2$ and $\gamma_1 = \gamma_2$, provided the solution of (1.1)–(1.3) exists. But they only declared the existence and uniqueness of global small smooth solution under the assumption that the initial data were sufficiently small in H^s where $s > d/2 + 2$. And in [3], Castella proved the global existence and the asymptotic behavior of solutions in the function space L^2 for the mixed-state unipolar Schrödinger-Poisson systems without the defocusing nonlinearity. In [6], with the help of madelung transform and WKB expansion, Li and Lin discussed the following unipolar nonlinear Schrödinger-Poisson system:

$$\begin{aligned} i\varepsilon \psi_t^\varepsilon + \frac{\varepsilon^2}{2} \Delta \psi^\varepsilon - (V^\varepsilon(x, t) + f'(|\psi^\varepsilon|^2)) \psi^\varepsilon - (\arg \psi^\varepsilon) \psi^\varepsilon &= 0, \\ -\Delta V^\varepsilon &= |\psi^\varepsilon|^2 - C(x), \quad V \rightarrow 0 \text{ as } |x| \rightarrow \infty, \end{aligned}$$

subject to the rapidly oscillating (WKB) initial condition

$$\psi^\varepsilon(x, 0) = \psi_0^\varepsilon = A_0^\varepsilon(x) e^{\frac{i}{\varepsilon} S_0(x)}, \quad (1.5)$$

where $f \in C^\infty(\mathbb{R}^+; \mathbb{R})$, $S_0 \in H^s(\mathbb{R}^d)$, $d \geq 1$, for $s \geq d/2 + 2$, A_0^ε was a function, polynomial in ε , with coefficients of Sobolev regularity in x , and the function $C(x) > 0$ denoted the background ions. They obtained the existence of smooth solution where the wave function was of the form $\psi^\varepsilon(x, t) = A^\varepsilon(x, t) e^{\frac{i}{\varepsilon} S^\varepsilon(x, t)}$, with A^ε and ∇S^ε bounded in $L^\infty([0, T]; H^s(\mathbb{R}^d))$ and the initial data being sufficiently small in $H^s(\mathbb{R}^d)$. However, to our knowledge, there is no previous result on the global existence and the asymptotic behavior of solutions for the (BDNLSP) system with arbitrary initial data in $\Sigma(\mathbb{R}^3)$. In this paper, by using the pseudo-conformal conservation law of the (BDNLSP) system and applying the time-space $L^p - L^{p'}$ estimate method, we shall establish the global existence and uniqueness of the solution to the (BDNLSP) system with initial data in $\Sigma(\mathbb{R}^3)$. As a byproduct, the large time behavior to the solution is also obtained. Although the above results are established for the single bipolar defocusing nonlinear Schrödinger-Poisson system, the results can be extended to the mixed-state bipolar defocusing nonlinear Schrödinger-Poisson system within the same framework.

For convenience, we first introduce some notation. For any $p \in [2, \infty)$, we denote $\frac{1}{\gamma(p)} = \frac{3}{2}(\frac{1}{2} - \frac{1}{p})$. $S(t)$ denotes the unitary group generated by $\frac{\varepsilon}{2}i\Delta$ in $L^2(\mathbb{R}^3)$. For $p \in [1, \infty]$, we denote by p' the conjugate exponent of p , defined by $1/p + 1/p' = 1$. \bar{z} denotes the conjugate of the complex number z .

Now we state the main result of this paper.

THEOREM 1.1 (Existence and uniqueness). Let $\varphi_j \in \Sigma(\mathbb{R}^3)$. Assume that $\rho \in [2, 6)$. Then, there exists a unique solution

$$\psi_j \in \mathcal{C}(\mathbb{R}; \Sigma(\mathbb{R}^3)) \cap L^\infty(\mathbb{R}; H^1(\mathbb{R}^3)) \cap L_{loc}^{\gamma(\rho)}(\mathbb{R}; H_\rho^1(\mathbb{R}^3)), \text{ for } j = 1, 2$$

to the (BDNLSP) system with the initial data (1.3).

Moreover, the solution (ψ_1, ψ_2, V) satisfies the L^2 -norm, the energy, and the pseudo-conformal conservation laws (for details, one can see Proposition 2.1 in the case $d = 3$).

THEOREM 1.2 (Large time behavior). Let (ψ_1, ψ_2, V) and ρ be as in Theorem 1.1. Then, there exist constants C depending only on $\|\varphi_j\|_{H^1}$ and $\|x|\varphi_j\|_2$ such that

$$\|\psi_j\|_\rho \leq C|t|^{-\frac{1}{\gamma(\rho)}}, \quad \forall \rho \in [2, 6), \quad \forall |t| \geq 1, \quad (1.6)$$

$$\|\nabla V(t)\|_\rho \leq C|t|^{-(1-\frac{3}{2\rho})}, \quad \forall \rho \in (\frac{3}{2}, \infty), \quad \forall |t| \geq 1, \quad (1.7)$$

$$\|V(t)\|_\rho \leq C|t|^{-\frac{1}{2}(1-\frac{3}{\rho})}, \quad \forall \rho \in (3, \infty), \quad \forall |t| \geq 1. \quad (1.8)$$

In the next section we shall derive three conservation identities including the L^2 -norm, the energy, and the pseudo-conformal conservation laws in the whole space \mathbb{R}^d for any $d \in \mathbb{N}$. In Sec. 3, we will give some basic estimates used in our proofs for $d = 3$. Section 4 is devoted to the proof of the existence and uniqueness of the solutions for the initial data in $\Sigma(\mathbb{R}^3)$. Finally, the large time behavior of the solution is obtained in Sec. 5.

2. Derivation of the conservation laws.

PROPOSITION 2.1. Let $d \in \mathbb{N}$, $\{\psi_j\}$ be a solution of the (BDNLSP) system with the initial value $\varphi_j(x) \in \Sigma(\mathbb{R}^d)$. Then, we have the following conservation laws for all $t \in \mathbb{R}$:

(i) L^2 -norm law:

$$\|\psi_j(t)\|_2 = \|\varphi_j\|_2 \quad \text{for } j = 1, 2; \quad (2.1)$$

(ii) Energy conservation law:

$$\varepsilon^2 \sum_{j=1}^2 \|\nabla \psi_j(t)\|_2^2 + \lambda^2 \|\nabla V\|_2^2 + 2 \sum_{j=1}^2 \frac{a_j^2}{\gamma_j + 1} \|\psi_j(t)\|_{2(\gamma_j+1)}^{2(\gamma_j+1)} = \text{const}; \quad (2.2)$$

(iii) Pseudo-conformal conservation law (cf. [9]):

$$\begin{aligned} & \sum_{j=1}^2 \|x\psi_j + i\varepsilon t \nabla \psi_j\|_2^2 + \lambda^2 t^2 \|\nabla V\|_2^2 + 2t^2 \sum_{j=1}^2 \frac{a_j^2}{\gamma_j + 1} \|\psi_j\|_{2(\gamma_j+1)}^{2(\gamma_j+1)} \\ & \quad + 2 \sum_{j=1}^2 \frac{a_j^2(d\gamma_j - 2)}{\gamma_j + 1} \int_0^t \tau \|\psi_j(\tau)\|_{2(\gamma_j+1)}^{2(\gamma_j+1)} d\tau \\ & = \sum_{j=1}^2 \|x|\varphi_j\|_2^2 + (4-d)\lambda^2 \int_0^t \tau \|\nabla V(\tau)\|_2^2 d\tau. \end{aligned} \quad (2.3)$$

Proof. Denote $eq(\psi_j) =: i\varepsilon\dot{\psi}_j + \frac{\varepsilon^2}{2}\Delta\psi_j - (q_jV + h_j(|\psi_j|^2))\psi_j$.

(i) It is well known that (2.1) holds for $j = 1, 2$. We omit its proof.

(ii) We consider

$$\Re(eq(\psi_j), \dot{\psi}_j) = 0$$

where (\cdot, \cdot) denotes the L^2 -inner product. From the above, we can get

$$\int_{\mathbb{R}^d} \left\{ \frac{\varepsilon}{4} \partial_t |\nabla \psi_j|^2 + \frac{1}{2} (q_j V + h_j(|\psi_j|^2)) \partial_t |\psi_j|^2 \right\} dx = 0.$$

Hence,

$$\begin{aligned} \int_{\mathbb{R}^d} \left\{ \frac{\varepsilon}{4} \partial_t (|\nabla \psi_1|^2 + |\nabla \psi_2|^2) + \frac{1}{2} V \partial_t (|\psi_1|^2 - |\psi_2|^2) \right. \\ \left. + \frac{1}{2} h_1(|\psi_1|^2) \partial_t |\psi_1|^2 + \frac{1}{2} h_2(|\psi_2|^2) \partial_t |\psi_2|^2 \right\} dx = 0. \end{aligned} \quad (2.4)$$

Integrating (2.4) over $[0, t]$, we obtain the desired identity.

(iii) Considering

$$\Re(eq(\psi_j), \bar{\psi}_j) = 0,$$

we have

$$-\varepsilon \Im \partial_t \psi_j \bar{\psi}_j + \frac{\varepsilon^2}{2} \Re \nabla \cdot (\nabla \psi_j \bar{\psi}_j) - \frac{\varepsilon^2}{2} |\nabla \psi_j|^2 - (q_j V + h_j(|\psi_j|^2)) |\psi_j|^2 = 0. \quad (2.5)$$

Noticing

$$\Re(eq(\psi_j), \bar{\psi}_{j,r}) = 0 \quad \text{with } r := |x|,$$

we obtain

$$\begin{aligned} -\varepsilon \Im \partial_t \psi_j \bar{\psi}_{j,r} + \frac{\varepsilon^2}{2} \Re \nabla \cdot (\nabla \psi_j \bar{\psi}_{j,r}) - \frac{r}{2} |\nabla \psi_j|^2 + \frac{\varepsilon^2}{4} (d-2) |\nabla \psi_j|^2 \\ - \frac{1}{2} (q_j V + h_j(|\psi_j|^2)) r \partial_r |\psi_j|^2 = 0. \end{aligned} \quad (2.6)$$

Due to

$$\partial_t (\psi_j \bar{\psi}_{j,r}) + \nabla \cdot (x \bar{\psi}_j \partial_t \psi_j) = 2 \partial_t \psi_j \bar{\psi}_{j,r} + 2 \Re \psi_j x \cdot \nabla \partial_t \bar{\psi}_j + d \partial_t \psi_j \bar{\psi}_j,$$

we have, by taking the imaginary part, that

$$\Im \partial_t (\psi_j \bar{\psi}_{j,r}) + \Im \nabla \cdot (x \bar{\psi}_j \partial_t \psi_j) = d \Im \partial_t \psi_j \bar{\psi}_j + 2 \Im (\partial_t \psi_j \bar{\psi}_{j,r}).$$

From (2.5) and (2.6), we have

$$\begin{aligned} -\varepsilon \Im \partial_t (\psi_j \bar{\psi}_{j,r}) - \Im \nabla \cdot (x \bar{\psi}_j \partial_t \psi_j) - \frac{\varepsilon^2}{2} d \Re \nabla \cdot (\nabla \psi_j \bar{\psi}_j) - d (q_j V + h_j(|\psi_j|^2)) |\psi_j|^2 \\ + \varepsilon^2 \Re \nabla \cdot (\nabla \psi_j \bar{\psi}_{j,r}) - \frac{r}{2} |\nabla \psi_j|^2 - (q_j V + h_j(|\psi_j|^2)) r \partial_r |\psi_j|^2 = 0. \end{aligned}$$

We also have

$$\partial_t |x \psi_j + i \varepsilon t \nabla \psi_j|^2 = \partial_t (|\psi_j r|^2 + \varepsilon^2 t^2 |\nabla \psi_j|^2 + 2 \varepsilon t \Im \psi_j x \cdot \nabla \bar{\psi}_j).$$

Since

$$\begin{aligned}
 \frac{\varepsilon}{2} \partial_t |\psi_j r|^2 &= \varepsilon \Re(\partial_t \psi_j \bar{\psi}_j) r^2 = \Im i \varepsilon \partial_t \psi_j \bar{\psi}_j r^2 \\
 &= \Im \left[-\frac{\varepsilon^2}{2} \Delta \psi_j + (q_j V + h_j(|\psi_j|^2)) \psi_j \right] \bar{\psi}_j r^2 \\
 &= -\frac{\varepsilon^2}{2} \Im \Delta \psi_j \bar{\psi}_j r^2, \\
 \Im \nabla \cdot (\nabla \psi_j \bar{\psi}_j r^2) &= \Im \Delta \psi_j \bar{\psi}_j r^2 - 2 \Im (\nabla \bar{\psi}_j \cdot x) \psi_j,
 \end{aligned}$$

we have

$$\begin{aligned}
 \partial_t |x \psi_j + i \varepsilon t \nabla \psi_j|^2 &= 2 \varepsilon^2 t |\nabla \psi_j|^2 + \varepsilon^2 t^2 \partial_t |\nabla \psi_j|^2 - \varepsilon \Im \nabla \cdot (\nabla \psi_j \bar{\psi}_j r^2) \\
 &\quad + 2 \varepsilon t \Im \partial_t (\psi_j x \cdot \nabla \bar{\psi}_j).
 \end{aligned} \tag{2.7}$$

Integrating (2.7) over \mathbb{R}^d , we obtain that for $j = 1, 2$

$$\begin{aligned}
 \partial_t \|x \psi_j + i \varepsilon t \nabla \psi_j\|_2^2 &= \varepsilon^2 t^2 \partial_t \|\nabla \psi_j\|_2^2 - 2t \int_{\mathbb{R}^d} \{d(q_j V + h_j(|\psi_j|^2)) |\psi_j|^2 \\
 &\quad + (q_j V + h_j(|\psi_j|^2)) r \partial_r |\psi_j|^2\} dx.
 \end{aligned} \tag{2.8}$$

From the above, we obtain

$$\begin{aligned}
 \partial_t \sum_{j=1}^2 \|x \psi_j + i \varepsilon t \nabla \psi_j\|_2^2 &+ 4t^2 \partial_t \left(\frac{\lambda^2}{4} \|\nabla V\|_2^2 + \sum_{j=1}^2 \frac{a_j^2}{2(\gamma_j + 1)} \|\psi_j\|_{2(\gamma_j + 1)}^{2(\gamma_j + 1)} \right) \\
 &+ 2t \int_{\mathbb{R}^d} \{dV(-\lambda^2 \Delta V) + V \partial_r(-\lambda^2 \Delta V) r + d \sum_{j=1}^2 a_j^2 |\psi_j|^{2(\gamma_j + 1)} \\
 &+ \sum_{j=1}^2 h_j(|\psi_j|^2) r \partial_r |\psi_j|^2\} dx = 0.
 \end{aligned} \tag{2.9}$$

Noticing that

$$h_j(s) r \partial_r s = a_j^2 s^{\gamma_j} r \partial_r s = \frac{a_j^2}{\gamma_j + 1} [\nabla \cdot (x s^{\gamma_j + 1}) - d s^{\gamma_j + 1}] \tag{2.10}$$

and

$$\begin{aligned}
 \int_{\mathbb{R}^d} V r \partial_r (-\Delta V) dx &= \int_{\mathbb{R}^d} \nabla V_r \cdot \nabla (V r) dx = \int_{\mathbb{R}^d} (\nabla V_r \cdot \nabla V) r + (\nabla V_r \cdot \nabla r) V dx \\
 &= \int_{\mathbb{R}^d} \frac{1}{2} r \partial_r |\nabla V|^2 dx + \int_{\mathbb{R}^d} V \Delta V dx \\
 &= \int_{\mathbb{R}^d} \frac{1}{2} [\nabla \cdot (x |\nabla V|^2) - d |\nabla V|^2] dx - \int_{\mathbb{R}^d} \nabla V \cdot \nabla V dx \\
 &= -\left(\frac{d}{2} + 1\right) \|\nabla V\|_2^2,
 \end{aligned} \tag{2.11}$$

we can obtain, in view of (2.4) and (2.8)–(2.11), that

$$\begin{aligned} \partial_t \Big[\sum_{j=1}^2 \|x\psi_j + i\varepsilon t \nabla \psi_j\|_2^2 + \lambda^2 t^2 \|\nabla V\|_2^2 + \sum_{j=1}^2 \frac{2a_j^2}{\gamma_j + 1} t^2 \|\psi_j\|_{2(\gamma_j+1)}^{2(\gamma_j+1)} \Big] \\ + 2 \sum_{j=1}^2 \frac{a_j^2(d\gamma_j - 2)}{\gamma_j + 1} t \|\psi_j(t)\|_{2(\gamma_j+1)}^{2(\gamma_j+1)} - (4-d)\lambda^2 t \|\nabla V(t)\|_2^2 = 0, \end{aligned}$$

which yields the pseudo-conformal conservation law. \square

3. Basic estimates.

LEMMA 3.1 (Estimate I). Let $\frac{1}{\rho'} = \frac{2}{a} + \frac{1}{\rho} + \frac{1}{3} - 1$, $u, v \in L^{\gamma(a)}(0, T; H_a^1)$ and $w \in L^{\gamma(\rho)}(0, T; H_\rho^1)$. Then we have the estimate

$$\begin{aligned} \left\| \left(\frac{1}{r} * uv \right) w \right\|_{L^{\gamma(\rho)'}(0, T; H_{\rho'}^1)} &\leq CT^{1/2} \|u\|_{L^{\gamma(a)}(0, T; H_a^1)} \\ &\quad \cdot \|v\|_{L^{\gamma(a)}(0, T; H_a^1)} \|w\|_{L^{\gamma(\rho)}(0, T; H_\rho^1)}. \end{aligned} \quad (3.1)$$

Proof. By the known estimate as in [8], we have the following:

$$\begin{aligned} \left\| \left(\frac{1}{r} * uv \right) w \right\|_{H_{\rho'}^1} &\leq C \left\| \frac{1}{r} * uv \right\|_{L^p} \|w\|_{H_q^1} + C \left\| \frac{1}{r} * uv \right\|_{H_q^1} \|w\|_{L^p} \\ &\leq C \|u\|_{H_{2m}^1} \|v\|_{H_{2m}^1} \|w\|_{H_q^1} \end{aligned}$$

where $\frac{1}{\rho'} = \frac{1}{p} + \frac{1}{q}$, $\frac{1}{p} = \frac{1}{m} + \frac{1}{3} - 1$. Let $2m = a$, $q = \rho$, i.e., $\frac{1}{\rho'} = \frac{2}{a} + \frac{1}{\rho} + \frac{1}{3} - 1$. By the Sobolev embedding theorem (cf. [1]), we obtain

$$\left\| \left(\frac{1}{r} * uv \right) w \right\|_{H_{\rho'}^1} \leq C \|u\|_{H_a^1} \|v\|_{H_a^1} \|w\|_{H_\rho^1}.$$

Since

$$\frac{1}{\gamma(\rho)'} = 1 - \frac{3}{2} \left(\frac{1}{\rho'} - \frac{1}{2} \right) = \frac{1}{2} + \frac{2}{\gamma(a)} + \frac{1}{\gamma(\rho)},$$

we have the desired result. \square

LEMMA 3.2 (Estimate II). Let $\rho \in [2, 6)$; we have

$$\| |u|^p u \|_{L^{\gamma(\rho)'}(0, T; H_{\rho'}^1)} \leq CT^{1 - \frac{2}{\gamma(\rho)}} \|u\|_{L^\infty(0, T; H^1)}^p \|u\|_{L^{\gamma(\rho)}(0, T; H_\rho^1)}. \quad (3.2)$$

Proof. From the identity

$$\begin{aligned} \nabla(|u|^p u) &= \nabla(|u|^p)u + |u|^p \nabla u = \frac{p}{2} |u|^{p-2} (\nabla u \bar{u} + u \nabla \bar{u})u + |u|^p \nabla u \\ &= \left(\frac{p}{2} + 1 \right) |u|^p \nabla u + \frac{p}{2} |u|^{p-2} u^2 \nabla \bar{u}, \end{aligned}$$

we have, in view of $\frac{1}{\rho'} = \frac{p}{q} + \frac{1}{\rho}$, that

$$\begin{aligned} \|\nabla(|u|^p u)\|_{L_{\rho'}^1} &\leq C \| |u|^p \nabla u \|_{L_{\rho'}^1} + C \| |u|^{p-2} u^2 \nabla \bar{u} \|_{L_{\rho'}^1} \\ &\leq C \|u\|_{L_a^p}^p \|\nabla u\|_{L^\rho}. \end{aligned}$$

By the Sobolev embedding theorem, it yields

$$\| |u|^p u \|_{H_{\rho'}^1} \leq C \|u\|_{H^1}^p \|u\|_{H_\rho^1},$$

which implies the desired result in view of the Hölder inequality with respect to the time variable. \square

Now we introduce the Galilei-type operator

$$J(t) = x + i\varepsilon t \nabla. \quad (3.3)$$

Let $\mathcal{M}(t) = e^{\frac{ix|t|^2}{2\varepsilon}}$ and $w_j = \mathcal{M}(-t)\psi_j$; we easily see that

$$J(t) = S(t)xS(-t) = \mathcal{M}(t)(i\varepsilon t \nabla)\mathcal{M}(-t), \quad (3.4)$$

$$|w_j(t)| = |\psi_j(t)|, \quad |J(t)\psi_j(t)| = \varepsilon|t||\nabla w_j(t)|. \quad (3.5)$$

LEMMA 3.3 (Estimate III). It holds

$$\|J(|\psi_j|^p \psi_j)\|_{L^{\gamma(\rho)'}(0,T;L^{\rho'})} \leq CT^{1-\frac{2}{\gamma(\rho)}} \|\psi_j\|_{L^\infty(0,T;H^1)}^p \|J\psi_j\|_{L^{\gamma(\rho)}(0,T;L^\rho)}. \quad (3.6)$$

Proof. We have, in view of (3.5), that

$$\|J(|\psi_j|^p \psi_j)\|_{L^{\rho'}} = \varepsilon|t| \|\nabla(|w_j|^p w_j)\|_{L^{\rho'}}. \quad (3.7)$$

Since

$$\nabla(|w_j|^p w_j) = \nabla(|w_j|^p)w_j + |w_j|^p \nabla w_j,$$

we obtain from Hölder's inequality

$$\|\nabla(|w_j|^p w_j)\|_{L^{\rho'}} \leq C \|w_j\|_{H^1}^p \|\nabla w_j\|_{L^\rho}. \quad (3.8)$$

Then, by (3.5), (3.7), and (3.8), we see that

$$\|J(|\psi_j|^p \psi_j)\|_{L^{\rho'}} \leq C \|\psi_j\|_{H^1}^p \|J\psi_j\|_{L^\rho},$$

which implies the desired result (3.6) in view of the Hölder inequality with respect to the time variable t . \square

LEMMA 3.4 (Estimate IV). We have the estimate

$$\begin{aligned} \|J(V\psi_j)\|_{L^{\gamma(\rho)'}(0,T;L^{\rho'})} &\leq CT^{1/2} \|(\psi_1, \psi_2)\|_{L^{\gamma(a)}(0,T;L^a)} \\ &\quad \cdot \|(J\psi_1, J\psi_2)\|_{L^{\gamma(\rho)}(0,T;L^\rho)} \end{aligned} \quad (3.9)$$

where $\|(u, v)\|_X := \|u\|_X + \|v\|_X$.

Proof. Noticing that

$$\begin{aligned} \nabla(Vw_j) &= \nabla Vw_j + V\nabla w_j, \\ \nabla V &= C\nabla\left(\frac{1}{r} * (|w_1|^2 - |w_2|^2)\right) = \frac{C}{r} * \nabla(|w_1|^2 - |w_2|^2) \\ &= \frac{C}{r} * (w_1 \nabla \bar{w}_1 + \nabla w_1 \bar{w}_1 - w_2 \nabla \bar{w}_2 - \nabla w_2 \bar{w}_2), \end{aligned}$$

we have for $\frac{1}{\rho'} = \frac{1}{\rho} + \frac{2}{a} + \frac{1}{3} - 1$

$$\|\nabla(Vw_j)\|_{L^{\rho'}} \leq C(\|w_1\|_{L^a}^2 + \|w_2\|_{L^a}^2)(\|\nabla w_1\|_{L^\rho} + \|\nabla w_2\|_{L^\rho}).$$

Thus, we can get

$$\begin{aligned} \|J(V\psi_j)\|_{L^{\rho'}} &= \varepsilon|t| \|\nabla(Vw_j)\|_{L^{\rho'}} \leq C\varepsilon|t|(\|w_1\|_{L^a}^2 + \|w_2\|_{L^a}^2)(\|\nabla w_1\|_{L^\rho} + \|\nabla w_2\|_{L^\rho}) \\ &\leq C(\|\psi_1\|_{L^a}^2 + \|\psi_2\|_{L^a}^2)(\|J\psi_1\|_{L^\rho} + \|J\psi_2\|_{L^\rho}), \end{aligned}$$

which implies the desired result. \square

LEMMA 3.5 (Estimate V). We have the following estimate

$$\begin{aligned} & \| (V_k \psi_{1k} - V_l \psi_{1l}, V_k \psi_{2k} - V_l \psi_{2l}) \|_{L^{\gamma(\rho)'}(0,T; H_{\rho'}^1)} \\ & \leq CT^{1/2} \| (\psi_{1k}, \psi_{2k}, \psi_{1l}, \psi_{2l}) \|_{L^{\gamma(a)}(0,T; H_a^1)}^2 \\ & \quad \cdot \| (\psi_{1k} - \psi_{1l}, \psi_{2k} - \psi_{2l}) \|_{L^{\gamma(\rho)}(0,T; H_{\rho}^1)}. \end{aligned} \quad (3.10)$$

Proof. Since

$$\begin{aligned} & C(V(\psi_{1k}, \psi_{2k})\psi_{jk} - V(\psi_{1l}, \psi_{2l})\psi_{jl}) \\ & = \left(\frac{1}{r} * (|\psi_{1k}|^2 - |\psi_{2k}|^2)\right)\psi_{jk} - \left(\frac{1}{r} * (|\psi_{1l}|^2 - |\psi_{2l}|^2)\right)\psi_{jl} \\ & = \left(\frac{1}{r} * (|\psi_{1k}|^2 - |\psi_{2k}|^2)\right)(\psi_{jk} - \psi_{jl}) + \psi_{jl} \left[\frac{1}{r} * ((|\psi_{1k}|^2 - |\psi_{1l}|^2) - (|\psi_{2k}|^2 - |\psi_{2l}|^2))\right] \\ & = \left(\frac{1}{r} * (|\psi_{1k}|^2 - |\psi_{2k}|^2)\right)(\psi_{jk} - \psi_{jl}) + \left[\frac{1}{r} * ((\bar{\psi}_{1k} - \bar{\psi}_{1l})\psi_{1k}) + \frac{1}{r} * ((\psi_{1k} - \psi_{1l})\bar{\psi}_{1l})\right] \\ & \quad + \frac{1}{r} * ((\bar{\psi}_{2k} - \bar{\psi}_{2l})\psi_{2k}) + \frac{1}{r} * ((\psi_{2k} - \psi_{2l})\bar{\psi}_{2l})\psi_{jl}, \end{aligned}$$

we have the desired result by the Hölder inequality. \square

4. The proof of the existence. In this section, we will prove the local existence of the Cauchy problem (BDNLSP) with the initial data (1.3) first. Let $S(t) := e^{\frac{1}{2}i\varepsilon\Delta t}$ and consider the integral equation

$$\psi_j(t) = S(t)\varphi_j - \frac{1}{\varepsilon}i \int_0^t S(t-\tau)(q_j V(\tau) + h_j(|\psi_j(\tau)|^2))\psi_j(\tau) d\tau. \quad (4.1)$$

Define the workspace (\mathcal{D}, d) as

$$\mathcal{D} := \{(\psi_j)_{j=1,2} : \| \psi_j \|_{L^\infty(0,T; H^1) \cap L^{\gamma(a)}(0,T; H_a^1)} \leq M, \text{ for any } a \in [2, 6)\}, \quad (4.2)$$

with the distance

$$d((\psi_{1k}, \psi_{2k}), (\psi_{1l}, \psi_{2l})) = \| (\psi_{1k} - \psi_{1l}, \psi_{2k} - \psi_{2l}) \|_{L^{\gamma(\rho)}(0,T; H_{\rho}^1)} \quad (4.3)$$

where $M \geq 2 \max_{j=1,2} \|\varphi_j\|_{H^1}$ and $\rho \in [2, 6)$. It is clear that (\mathcal{D}, d) is a Banach space. Let us consider the mapping $\mathcal{T} = \mathcal{T}_1 \otimes \mathcal{T}_2 : (\mathcal{D}, d) \rightarrow (\mathcal{D}, d)$ defined by

$$\mathcal{T}_j : \psi_j(t) \mapsto S(t)\varphi_j - \frac{1}{\varepsilon}i \int_0^t S(t-\tau)(q_j V(\tau) + h_j(|\psi_j(\tau)|^2))\psi_j(\tau) d\tau, \quad j = 1, 2. \quad (4.4)$$

By Lemmas 3.1–3.5 and the Strichartz estimates (cf. [5]), we have

$$\begin{aligned}
 \|\mathcal{T}_j \psi_j\|_{L^{\gamma(\rho)}(0,T;H_\rho^1)} &\leq \|\varphi_j\|_{H^1} + \|V\psi_j\|_{L^{\gamma(\rho)'}(0,T;H_{\rho'}^1)} + \|h_j(|\psi_j|^2)\psi_j\|_{L^{\gamma(\rho)'}(0,T;H_{\rho'}^1)} \\
 &\leq \|\varphi_j\|_{H^1} + CT^{1/2} \left(\sum_{j=1}^2 \|\psi_j\|_{L^{\gamma(\rho)}(0,T;H_\rho^1)}^2 \right) \|\psi_j\|_{L^{\gamma(\rho)}(0,T;H_\rho^1)} \\
 &\quad + CT^{1-2/\gamma(\rho)} \|\psi_j\|_{L^\infty(0,T;H^1)}^{2\gamma_j} \|\psi_j\|_{L^{\gamma(\rho)}(0,T;H_\rho^1)} \\
 &\leq M/2 + (CT^{1/2}M^2 + CT^{1-2/\gamma(\rho)}M^{2\gamma_j})M \\
 &\leq M,
 \end{aligned} \tag{4.5}$$

where we have taken T so small that $CT^{1/2}M^2 + CT^{1-2/\gamma(\rho)}M^{2\gamma_j} < \frac{1}{2}$. Similar to the above, a straightforward computation shows that it holds

$$\begin{aligned}
 &\|\mathcal{T}(\psi_{1k}, \psi_{2k}) - \mathcal{T}(\psi_{1l}, \psi_{2l})\|_{L^{\gamma(\rho)}(0,T;H_\rho^1)} \\
 &\leq \frac{1}{2} \|(\psi_{1k} - \psi_{1l}, \psi_{2k} - \psi_{2l})\|_{L^{\gamma(\rho)}(0,T;H_\rho^1)}.
 \end{aligned} \tag{4.6}$$

Hence, \mathcal{T} is a contracted mapping from the Banach space (\mathcal{D}, d) to itself. By the Banach contraction mapping principle, we know that there exists a unique solution $(\psi_1, \psi_2) \in L^{\gamma(\rho)}(0, T; H_\rho^1) \times L^{\gamma(\rho)}(0, T; H_\rho^1)$ to the (BDNLSP) system with the initial data (1.3). From (4.1), (3.4) and (3.5), we may easily obtain $J\psi_1, J\psi_2 \in L^{\gamma(\rho)}(0, T; L^\rho)$ with the help of Lemmas 3.3–3.4. Thus, we can use the standard argument (cf. [3]) to extend it to a global one satisfying for any $T > 0$

$$\psi_1(t, x), \psi_2(t, x) \in \mathcal{C}(\mathbb{R}; \Sigma(\mathbb{R}^3)) \cap L^\infty(\mathbb{R}; H^1(\mathbb{R}^3)) \cap L^{\gamma(\rho)}(-T, T; H_\rho^1(\mathbb{R}^3)),$$

and prove the uniqueness of the global solution. We omit the details.

5. Large time behavior of the solution. By the pseudo-conformal conservation law, we get for $d = 3$

$$\lambda^2 t^2 \|\nabla V\|_2^2 \leq I + \lambda^2 \int_1^t \tau \|\nabla V(\tau)\|_2^2 d\tau, \tag{5.1}$$

where $I := \sum_{j=1}^2 \|x|\psi_j\|_2^2 + \lambda^2 \int_0^1 \tau \|\nabla V(\tau)\|_2^2 d\tau$.

From the Gronwall inequality, we have

$$\|\nabla V\|_2 \leq \frac{I^{1/2}}{\lambda} |t|^{-1/2}. \tag{5.2}$$

By the energy conservation law and the Sobolev embedding theorem, we obtain

$$\begin{aligned} \lambda^2 \|\nabla V\|_2^2 &\leq \varepsilon^2 \sum_{j=1}^2 \|\nabla \varphi_j\|_2^2 + \|\nabla V(0)\|_2^2 + \sum_{j=1}^2 \frac{2a_j^2}{\gamma_j + 1} \|\varphi_j\|_{2(\gamma_j+1)}^{2(\gamma_j+1)} \\ &\leq C(\varepsilon, \gamma_j, \|\varphi_j\|_{H^1}) + \|\nabla V(0)\|_2^2. \end{aligned} \quad (5.3)$$

$$\begin{aligned} \|\nabla V(0)\|_2 &= C \left\| \nabla \left(\frac{1}{|x|} * (|\varphi_1|^2 - |\varphi_2|^2) \right) \right\|_2 \leq C \left\| \frac{1}{|x|^2} * (|\varphi_1|^2 - |\varphi_2|^2) \right\|_2 \\ &\leq C \sum_{j=1}^2 \|\varphi_j\|_{L^{12/5}}^2 \leq C \sum_{j=1}^2 \|\varphi_j\|_{H^1}^2 \\ &\leq C(\varepsilon, \gamma_j, \|\varphi_j\|_{H^1}). \end{aligned} \quad (5.4)$$

Therefore, we have the estimate

$$\|\nabla V\|_2 \leq \frac{C}{\lambda} |t|^{-\frac{1}{2}}. \quad (5.5)$$

By the Sobolev embedding theorem and the pseudo-conformal conservation law, we have

$$\begin{aligned} \|\psi_j\|_\rho &= \|\mathcal{M}(-t)\psi_j\|_\rho \leq C \|\nabla \mathcal{M}(-t)\psi_j\|_2^{2/\gamma(\rho)} \|\mathcal{M}(t)\psi_j\|_2^{1-\gamma(\rho)} \\ &\leq C \|\mathcal{M}(-t)\left(\frac{x}{i\varepsilon t}\psi_j + \nabla \psi_j\right)\|_2^{2/\gamma(\rho)} \|\psi_j\|_2^{1-\gamma(\rho)} \\ &\leq C|t|^{-1/\gamma(\rho)}. \end{aligned} \quad (5.6)$$

From the above and the Hardy-Littlewood-Sobolev inequality, we obtain

$$\|\nabla V(t)\|_\rho \leq C|t|^{-(1-\frac{3}{2\rho})}, \quad \forall \rho \in \left(\frac{3}{2}, \infty\right), \quad \forall |t| \geq 1, \quad (5.7)$$

$$\|V(t)\|_\rho \leq C|t|^{-\frac{1}{2}(1-\frac{3}{\rho})}, \quad \forall \rho \in (3, \infty), \quad \forall |t| \geq 1. \quad (5.8)$$

REFERENCES

- [1] J. Bergh and J. Löfström, *Interpolation spaces, An introduction*, Grundlehren der Mathematischen Wissenschaften, No. 223, Springer-Verlag: Berlin-New York, 1976
- [2] F. Brezzi and P. A. Markowich, *The three-dimensional Wigner-Poisson problem: existence, uniqueness and approximation*, Math. Meth. Appl. Sci., **14**, 35-62 (1991)
- [3] F. Castella, *L^2 solutions to the Schrödinger-Poisson system: existence, uniqueness, time behaviour, and smoothing effects*, Math. Models Methods Appl. Sci., **7**(8), 1051-1083 (1997)
- [4] A. Jüngel and S. Wang, *Convergence of nonlinear Schrödinger-Poisson systems to the compressible Euler equations*, Comm. Part. Diff. Eqs. **28**, 1005-1022 (2003)
- [5] M. Keel and T. Tao, *Endpoint Strichartz estimates*, Amer. J. Math., **120**, 955-980 (1998)
- [6] H. L. Li and C. K. Lin, *Semiclassical limit and well-posedness of nonlinear Schrödinger-Poisson systems*, Electron. J. Diff. Eqns., **2003**, No. 93, pp. 1-17 (2003)
- [7] P. A. Markowich, G. Rein and G. Wolansky, *Existence and nonlinear stability of stationary states of the Schrödinger-Poisson system*, J. Statist. Phys., **106**(5-6), 1221-1239 (2002)
- [8] M. E. Taylor, *Tools for PDE, Pseudodifferential operators, paradifferential operators, and layer potentials*, Mathematical Surveys and Monographs, 81, American Mathematical Society, Providence, RI, 2000
- [9] B. X. Wang, *Large time behavior of solutions for critical and subcritical complex Ginzburg-Landau equations in H^1* , Science in China (Series A), **46**(1), 64-74 (2003)