

# TRANSFORMATIONS PRESERVING THE GRASSMANNIAN

BY  
WILLIAM C. NEMITZ

1. **Introduction.** For  $m$  a positive integer, let  $E_m$  be the arithmetic  $m$ -space over a commutative field  $F$ . Let  $\mathcal{A}_m$  be the full linear group of  $E_m$ , and let  $S_{m-1}$  be the projective space of homogeneous coordinates in  $E_m$ . For the rest of the paper, we fix two positive integers  $n$  and  $k$ , such that  $k < n$ . Let  $N = \binom{n}{k}$ , and let  $\Omega(k, n)$  be the  $k, n$  Grassmannian variety:

$$\Omega(k, n) \subset S_{N-1}.$$

Let  $\psi(k, n)$  be the set of those nonzero elements  $x$  of  $E_N$  such that there is some  $y$  satisfying

$$x \in y \in \Omega(k, n).$$

Let  $G$  be the set of nonsingular linear transformations of  $E_N$  which keep  $\psi(k, n)$  fixed as a set. If  $C_N$  is the center of the full linear group of  $E_N$ , then  $G/C_N$  is the set of projective transformations of  $S_{N-1}$  which keep  $\Omega(k, n)$  fixed as a set.

Let  $A(n, k)$  be the group of all  $k$ -compounds [1, Vol. 1, p. 291] of elements of  $\mathcal{A}_n$ . Then  $A(n, k)/(C_N \cap A(n, k))$  may be thought of as the group of projective transformations of  $S_{N-1}$  "induced" by the group of projective transformations of  $S_{n-1}$ . Since  $A(n, k)/(C_N \cap A(n, k))$  is isomorphic to  $(A(n, k) \cdot C_N)/C_N$ , and since  $A(n, k) \cdot C_N$  is a subgroup of  $G$ ,  $(A(n, k) \cdot C_N)/C_N$  is a subgroup of  $G/C_N$ .

The principal results to be proved here are:

1. If  $n \neq 2k$ , then

$$A(n, k) \cdot C_N = G,$$

and thus

$$(A(n, k) \cdot C_N)/C_N = G/C_N.$$

2. If  $n = 2k$ , let  $J$  denote the "star dual" mapping of  $\psi(k, n)$  onto itself (see 2). Since

$$J^2 = (-1)^{\binom{k^2}{2}} I,$$

where  $I$  is the identity element of  $\mathcal{A}_N$ ,  $J$  generates a cyclic subgroup of order 2 if  $k$  is even, and of order 4 if  $k$  is odd. Let  $\mathcal{J}$  denote this group. Let  $\mathcal{X}$  be the

---

Received by the editors May 28, 1962 and, in revised form, December 12, 1962.

subgroup of  $G/C_N$  made up of cosets of elements of  $\mathcal{J}$ . Thus  $\mathcal{K}$  is of order 2. Then, in this case,

$$\mathcal{J} \cdot A(n, k) \cdot C_N = G,$$

and thus

$$\mathcal{K} \cdot ((A(n, k) \cdot C_N)/C_N) = G/C_N.$$

2. **Notation.** (For definitions of terms used here and proofs of results given here, see [2].) We shall denote the exterior product of vectors by “ $\wedge$ ”. Thus  $x$  is an element of  $\psi(k, n)$  if and only if there is a linearly independent set of  $k$  elements of  $E_n, x_1, x_2, x_3, \dots, x_k$ ; and

$$x = x_1 \wedge x_2 \wedge x_3 \wedge \dots \wedge x_k.$$

For  $A \in \mathcal{A}_n$ , let  $A^k$  be the  $k$ -compound of  $A$ . Thus if

$$x = x_1 \wedge x_2 \wedge x_3 \wedge \dots \wedge x_k,$$

then

$$A^k x = Ax_1 \wedge Ax_2 \wedge Ax_3 \wedge \dots \wedge Ax_k.$$

For  $E \subset E_m$ , let  $L(E)$  be the subspace of  $E_m$  spanned by  $E$ . If  $x \in \psi(k, n)$ , such that

$$x = x_1 \wedge x_2 \wedge x_3 \wedge \dots \wedge x_k,$$

let

$$\pi(x) = L(\{x_1, x_2, x_3, \dots, x_k\}).$$

For any positive integer  $m$ , let

$$\mathcal{N}(m) = \{1, 2, 3, \dots, m\}.$$

For  $t$  a positive integer,  $t \leq m$ , let

$$P(m, t) = \{p: p = \{p_1, p_2, p_3, \dots, p_t\}, p_i \in \mathcal{N}(m) \text{ for } i \in \mathcal{N}(t), \text{ and}$$

$$p_1 < p_2 < p_3 < \dots < p_t\}.$$

For  $p \in P(m, t)$ , let  $c(p)$  be that element of  $P(m, m - t)$  such that

$$p \cup c(p) = \mathcal{N}(m).$$

For  $x$  an element of  $\psi(k, n)$ ,  $*x$  is that element of  $\psi(n - k, n)$  defined by

$$(*x)_q = \varepsilon(q) x_{c(q)},$$

where  $q$  is any element of  $P(n, n - k)$ , and  $\varepsilon(q)$  is  $-1$  to the power of the parity of the permutation  $(q_1, q_2, q_3, \dots, q_{n-k}, (cq)_1, (cq)_2, (cq)_3, \dots, (cq)_k)$ . Let  $J$  be that mapping of  $\psi(k, n)$  onto  $\psi(n - k, n)$  defined by

$$J(x) = *x.$$

Then  $J$  can be extended to a nonsingular linear mapping of  $E_N$  onto itself.

Since  $k < n$ , we may consider  $E_{k+1}$  as a subspace of  $E_n$ , and  $\psi(k, k + 1)$  as a subset of  $\psi(k, n)$ . On occasion, we shall find it necessary to use the  $*$ -dual of a vector in  $\psi(k, k + 1)$  "relative to  $E_{k+1}$ ." That is, for  $x$  an element of  $\psi(k, k + 1) \subset \psi(k, n)$ ,

$$(*_{k+1}x)_i = (-1)^{i-1}x_{c(i)}, \quad \text{where } c(i) = \mathcal{N}(k + 1) - \{i\}, \text{ if } 1 \leq i \leq k + 1;$$

and

$$(*_{k+1}x)_i = 0, \quad \text{if } i > k + 1.$$

Then  $*_{k+1}x \in E_{k+1} \subset E_n$ , and

$$L(*_{k+1}x) = (\pi(x))^{\perp_{k+1}},$$

where  $\perp_{k+1}$  denotes the orthogonal complement relative to  $E_{k+1}$ .

For  $i \in \mathcal{N}(m)$ , let  $e_i$  be that element of  $E_m$  whose  $j$ th component is  $\delta_{ij}$ . For  $p \in P(n, k)$ , let

$$e_p = e_{p_1} \wedge e_{p_2} \wedge e_{p_3} \wedge \cdots \wedge e_{p_k}.$$

Then the set  $\{e_p : p \in P(n, k)\}$  is a basis for  $E_N$ .

For  $A \in G$ , and  $p \in P(n, k)$ , let  $A_p = Ae_p$ . Then  $A_p \in E_N$ , and it is the  $p$ th column vector of the matrix of  $A$ . For any  $q \in P(n, k - 1)$ ,

$$\dim\left(\bigcap \pi(e_p)\right) = k - 1,$$

the intersection being taken over all  $p \in P(n, k)$  such that  $q \subset p$ ; and

$$\dim(L(\{e_p : q \subset p \in P(n, k)\})) = n - k + 1.$$

So if  $A \in G$ , and  $q \in P(n, k - 1)$ , and if

$$M = A(L(\{e_p : q \subset p \in P(n, k)\})),$$

then  $\dim M = n - k + 1$ , and  $M$  is spanned by the set  $\{A_p : q \subset p \in P(n, k)\}$ . Furthermore, for  $p \in P(n, k)$ ,  $A_p \in M$  if and only if  $q \subset p$ .

Since we have excluded the zero vector from  $\psi(k, n)$ , no linear subspace of  $E_N$  is contained in  $\psi(k, n)$ . However, if  $M$  is a linear subspace of  $E_N$ , we shall say  $M \subset \psi(k, n)$  if and only if for  $x \in M$ , if  $x \neq 0$ , then  $x \in \psi(k, n)$ .

**3. Principal results.** The principal results may now be stated in the following two theorems.

**3.1. THEOREM.** *If  $n \neq 2k$ , and  $A \in G$ , then there exists  $C \in C_N$  and  $B \in \mathcal{A}_m$  such that*

$$A = CB^k.$$

**3.2. THEOREM.** *If  $n = 2k$ , and if  $A \in G$ , then there exists  $C \in C_N$  and  $B \in \mathcal{A}_n$  such that either*

$$A = CB^k$$

or

$$A = CJB^k.$$

The proofs of these theorems depend on the following three lemmas, which will be proved in §§4 and 5.

3.3. LEMMA. For  $m$  an integer,  $2 \leq m \leq N$ , let  $M$  be a subspace of  $E_N$ , with  $\dim M = m$ , such that there exists a set  $\{x_1, x_2, x_3, \dots, x_m\} \subset \psi(k, n)$  and  $\{x_1, x_2, x_3, \dots, x_m\}$  spans  $M$ . Then,

1. if

$$\dim \bigcap_{i=1}^m \pi(x_i) = k - 1,$$

then  $M \subset \psi(k, n)$ ,

$$\dim \bigcap_{x \in M} \pi(x) = k - 1,$$

and

$$\dim L(\{\pi(x) : x \in M\}) = k + m - 1;$$

2. if

$$\dim L(\{\pi(x_i) : 1 \leq i \leq m\}) = k + 1,$$

then  $M \subset \psi(k, n)$ ,  $m \leq k + 1$ ,

$$\dim \bigcap_{x \in M} \pi(x) = k - m + 1,$$

and

$$\dim L(\{\pi(x) : x \in M\}) = k + 1.$$

In either case,  $M$  is the set of all  $k$ -vectors of  $k$  dimensional subspaces of  $E_n$  which contain  $\bigcap_{x \in M} \pi(x)$  and are contained in  $L(\{\pi(x) : x \in M\})$ .

3.4. LEMMA. For  $m$  an integer,  $2 \leq m \leq N$ , let  $M$  be a subspace of  $E_N$ , with  $\dim M = m$ , and assume that  $M \subset \psi(k, n)$ . Let  $\{x_1, x_2, x_3, \dots, x_m\}$  be any spanning set of  $M$ . Then either

$$\dim \bigcap_{i=1}^m \pi(x_i) = k - 1,$$

or

$$\dim L(\{\pi(x_i) : 1 \leq i \leq m\}) = k + 1.$$

3.5. LEMMA. If  $A \in G$ , and if, for each  $q \in P(n, k-1)$ ,

$$\dim \bigcap \pi(A_p) = k - 1,$$

the intersection being taken over all  $p$  such that

$$q \subset p \in P(n, k),$$

then there exists  $C \in C_N$  and  $B \in \mathcal{A}_n$  such that

$$A = CB^k.$$

**Proof of Theorem 3.1 assuming Lemmas 3.3, 3.4, and 3.5.** First assume that  $n > 2k$ . For  $q \in P(n, k-1)$ , let  $M(q)$  be the subspace of  $E_N$  spanned by the set  $\{A_p: q \subset p \in P(n, k)\}$ . Then  $M(q) \subset \psi(k, n)$ , and  $\dim M(q) = n - k + 1$ . But  $n - k + 1 > k + 1$ . So by 3.3 and 3.4,

$$\dim \bigcap \pi(A_p) = k - 1,$$

the intersection being taken over all  $p$  such that

$$q \subset p \in P(n, k).$$

The result follows from 3.5. Now assume that  $n < 2k$ . Then for  $x \in \psi(n-k, n)$ ,  $JAJ^{-1}(x) \in \psi(n-k, n)$ . Hence there exists  $C \in C_N$  and  $B \in \mathcal{A}_n$  such that

$$JAJ^{-1} = CB^{n-k}.$$

So

$$A = CJ^{-1}B^{n-k}J.$$

By the Laplace expansion of a determinant,

$$J^{-1}B^{n-k}J = (\det B)I(B^{-T})^k,$$

where  $-T$  denotes inverse transpose. Hence

$$A = C(\det B)I(B^{-T})^k.$$

This completes the proof.

**Proof of Theorem 3.2 assuming Lemmas 3.3, 3.4, and 3.5.** We first show that if

$$\dim L(\{\pi(A_p): q' \subset p \in P(n, k)\}) = k + 1,$$

for some  $q' \in P(n, k-1)$ , then

$$\dim L(\{\pi(A_p): q \subset p \in P(n, k)\}) = k + 1,$$

for every  $q \in P(n, k-1)$ . It suffices to consider  $q' = \{1, 2, 3, \dots, k-1\}$  and to assume that

$$\dim L(\{\pi(A_p): q' \subset p \in P(n, k)\}) = k + 1.$$

Select  $q \in P(n, k-1)$ , so ordered that if  $q_i \in q'$ , then  $q_i = i$ . Let  $q'' = \{2, 3, 4, \dots, k-1, q_1\}$ . We will show that

$$\dim L(\{\pi(A_p): q'' \subset p \in P(n, k)\}) = k + 1.$$

If  $q_1 = 1$ , there is nothing to prove. So assume that  $q_1 \neq 1$ . Let  $p'' = \{1, 2, 3, \dots, k-1, q_1\}$ , and let

$$M' = L(\{A_p: q' \subset p \in P(n, k)\}),$$

and

$$M'' = L(\{A_p: q'' \subset p \in P(n, k)\}).$$

Then

$$M' \cap M'' = L(A_{p''}),$$

so

$$\dim(M' \cap M'') = 1.$$

Now let  $Q' = L(\{\pi(A_p): q' \subset p \in P(n, k)\})$ , and  $Q'' = \bigcap \pi(A_p)$ , the intersection being taken over all  $p \in P(n, k)$  such that  $q'' \subset p$ , and assume that  $\dim Q'' = k-1$ . Then

$$Q'' \subset \pi(A_{p''}) \subset Q'.$$

So the set of all  $y \in \psi(k, n)$  such that  $Q'' \subset \pi(y) \subset Q'$  is a subspace of  $M' \cap M''$ , but by [1, Vol. 2, Chapter XIV, Theorem I], the dimension of this subspace is 2. So  $\dim(M' \cap M'') \geq 2$ . This is a contradiction. So by Lemma 3.4,

$$\dim L(\{\pi(A_p): q'' \subset p \in P(n, k)\}) = k + 1.$$

Continuing in this manner, working with one element of  $q$  at a time, we conclude that

$$\dim L(\{\pi(A_p): q \subset p \in P(n, k)\}) = k + 1.$$

Hence either  $A$  or  $JA$  satisfies the conditions of Lemma 3.5, so the result follows from the fact that  $J^2 = (-1)^{(k^2)}I$ .

**4. Linear subspaces contained in  $\psi(k, n)$ .** Lemmas 3.3 and 3.4 describe the linear subspaces of  $E_N$  which are contained in  $\psi(k, n)$  in the sense of 2. In this section we give proofs of these two lemmas.

**Proof of Lemma 3.3.** Select a set  $\{x_1, x_2, x_3, \dots, x_m\} \subset \psi(k, n)$ , such that  $\{x_1, x_2, x_3, \dots, x_m\}$  spans  $M$ , and assume that

$$\dim \bigcap_{i=1}^m \pi(x_i) = k - 1.$$

Then without loss of generality, we may assume that

$$x_i = e_1 \wedge e_2 \wedge e_3 \wedge \dots \wedge e_{k-1} \wedge e_{k+i-1}, \quad \text{for } i = 1, 2, 3, \dots, m.$$

Now let  $x \in M$ . Then there exist  $a_1, a_2, a_3, \dots, a_m$ , elements of  $F$ , such that  $x = \sum_{i=1}^m a_i x_i$ . So

$$x = e_1 \wedge e_2 \wedge e_3 \wedge \dots \wedge e_{k-1} \wedge \left( \sum_{i=1}^m a_i e_{k+i-1} \right).$$

Hence  $M \subset \psi(k, n)$ , and consists of those  $k$ -vectors of  $k$ -spaces containing  $L(\{e_1, e_2, e_3, \dots, e_{k-1}\})$ , and contained in  $L(\{e_1, e_2, e_3, \dots, e_{k+m-1}\})$ . Now assume that

$$\dim L(\{\pi(x_i): 1 \leq i \leq m\}) = k + 1.$$

Then without loss of generality, we may assume that

$$\pi(x_i) \subset L(\{e_1, e_2, e_3, \dots, e_{k+1}\})$$

for  $i = 1, 2, 3, \dots, m$ . Hence the  $x_i$  may be thought of as  $k$ -vectors in  $E_{k+1}$ . So if  $x \in M$ ,  $x = \sum_{i=1}^m a_i x_i$ , for suitable elements  $a_i$  of  $F$ , then  $x$  is a  $k$ -vector in  $E_{k+1}$ . Hence  $M \subset \psi(k, n)$ , and

$$\dim L(\{\pi(x): x \in M\}) = k + 1.$$

Also, the set  $\{*_k x_i: 1 \leq i \leq m\}$  spans an  $m$ -space of  $E_{k+1}$ , so  $m \leq k + 1$ , and since  $L(*_k x_i) = (\pi(x_i))^{\perp_{k+1}}$ ,

$$\dim \bigcap_{i=1}^m \pi(x_i) = k - m + 1.$$

But for  $x \in M$ ,  $L(*_k x) \subset L(\{*_k x_i: 1 \leq i \leq m\})$ , and so

$$\bigcap_{i=1}^m \pi(x_i) \subset \pi(x).$$

Hence

$$\dim \bigcap \pi(x) = k - m + 1,$$

the intersection being taken over all  $x \in M$ . This completes the proof.

**Proof of Lemma 3.4.** Since  $M \subset \psi(k, n)$ , the plane spanned by  $x_i$  and  $x_j$  lies in  $\psi(k, n)$ , for  $i \neq j$ ,  $i, j = 1, 2, 3, \dots, m$ . By [1, Vol. 2, Chapter XIV, Theorem I],

$$\dim(\pi(x_i) \cap \pi(x_j)) = k - 1.$$

So, without loss of generality, we may assume that

$$\pi(x_1) = L(\{e_1, e_2, e_3, \dots, e_k\}),$$

and

$$\pi(x_2) = L(\{e_2, e_3, e_4, \dots, e_{k+1}\}).$$

Now assume that there is some  $x_j$ , say  $x_3$ , such that

$$\pi(x_1) \cap \pi(x_2) \subset \pi(x_3).$$

Then we may assume that  $\pi(x_3) = L(\{e_2, e_3, e_4, \dots, e_k, e_{k+2}\})$ . Now assume that there is some  $x_i$ , such that  $\pi(x_i)$  does not contain  $\pi(x_1) \cap \pi(x_2)$ . Since

$$\dim(\pi(x_i) \cap \pi(x_1)) = \dim(\pi(x_i) \cap \pi(x_2)) = k - 1,$$

we can choose a spanning set  $\{u_1, u_2, u_3, \dots, u_k\}$  for  $\pi(x_i)$  such that  $u_i \in \pi(x_1)$ , for  $i = 1, 2, 3, \dots, k-1$ , and  $u_k \in \pi(x_2)$ . Hence

$$\pi(x_i) \subset L(\{e_1, e_2, e_3, \dots, e_{k+1}\}),$$

and so

$$\dim(\pi(x_i) \cap \pi(x_3)) < k-1.$$

But this contradicts the fact that  $\dim(\pi(x_i) \cap \pi(x_3)) = k-1$ . So

$$L(\{e_2, e_3, e_4, \dots, e_k\}) \subset \pi(x_i),$$

and hence

$$\dim \bigcap_{i=1}^m \pi(x_i) = k-1.$$

Thus far, we have shown that if any three of the spaces  $\pi(x_1), \pi(x_2), \pi(x_3), \dots, \pi(x_m)$  intersect in a  $k-1$  space, then they all intersect in a  $k-1$  space. Now assume that no three of these spaces intersect in a  $k-1$  space. Hence, for  $i \neq 1, 2$ ,  $\pi(x_i)$  does not contain  $\pi(x_1) \cap \pi(x_2)$ . So, as before,  $\pi(x_i) \subset L(\{e_1, e_2, e_3, \dots, e_{k+1}\})$ , and so

$$\dim L(\{\pi(x_i): 1 \leq i \leq m\}) = k+1.$$

**5. Proof of Lemma 3.5.** The proof is in two parts.

**PART 1.** We first prove that, given the assumptions of the lemma, there is a set  $\{x_1, x_2, x_3, \dots, x_n\} \subset E_n$ , such that

$$(1) \quad \pi(A_p) = L(\{x_{p_1}, x_{p_2}, x_{p_3}, \dots, x_{p_k}\})$$

for any  $p \in P(n, k)$ . The proof is by induction on the number of vectors which can be found satisfying (1). First note that the assumption that for any  $q \in P(n, k-1)$ , the dimension of the intersection of the spaces  $\pi(A_p)$  for  $q \subset p \in P(n, k)$  is  $k-1$ , implies that to each  $q \in P(n, k-1)$  there is assigned in a one-to-one manner, a  $k-1$  space  $S(q)$  of  $E_n$ , such that

$$S(q) = \pi(A_p) \cap \pi(A_r),$$

for any  $p \in P(n, k)$ , and  $r \in P(n, k)$ , such that  $p \neq r$ , and  $q \subset p \cap r$ . Obviously, there is a set  $\{x_1, x_2, x_3, \dots, x_k\} \subset E_n$  such that if  $p = \{1, 2, 3, \dots, k\}$ , then (1) is true. So, assume that there exists a set  $\{x_1, x_2, x_3, \dots, x_t\} \subset E_n$ , for some integer  $t$ ,  $k \leq t \leq n-1$ , such that (1) holds for any  $p \in P(t, k)$ . Let  $p = \{1, 2, 3, \dots, k-1, t+1\}$ . Then there exists an  $x_{t+1} \in E_n$  such that (1) holds for this  $p$ . Let  $q$  be an element of  $P(t, k-1)$ , so ordered that if  $q_s \in p$ , then  $q_s = s$ . Let  $\bar{p} = q \cup \{t+1\}$ . We wish to show that (1) holds for  $\bar{p}$ . We now define a family of elements of  $P(n, k)$  as follows:



$$p(0) = p,$$

$$p(j) = (p(j-1) - \{j\}) \cup \{q_j\},$$

for  $j = 1, 2, 3, \dots, k-1$ . We will show by induction on  $j$ , that (1) holds for each  $p(j)$ . This will complete the induction on  $t$ , since  $\bar{p} = p(k-1)$ . Obviously, (1) is true if  $j = 0$ . Assume that, for some  $j$ ,  $0 \leq j < k-1$ , (1) holds for  $p(j)$ . If  $q_{j+1} = j+1$ , then (1) holds for  $p(j+1)$ . So assume that  $q_{j+1} \notin p$ . We also assume that  $q_{j+1} \neq k$ . Let

$$p' = (p(j+1) - \{t+1\}) \cup \{k\},$$

$$r = (p(j+1) - \{t+1\}) \cup \{j+1\},$$

$$Z(i) = L(\{x_1, x_2, x_3, \dots, x_i\}),$$

for  $i$  equal  $t$  or  $t+1$ . Then

$$\pi(A_{p(j+1)}) \cap Z(t) = \pi(A_{p'}) \cap \pi(A_{p(j+1)}) = S(p' \cap p(j+1))$$

$$= \pi(A_{p'}) \cap \pi(A_r) = L(\{x_{p(j+1)_1}, x_{p(j+1)_2}, x_{p(j+1)_3}, \dots, x_{p(j+1)_{k-1}}\}),$$

and

$$\pi(A_{p(j)}) \cap \pi(A_{p(j+1)}) = S(p(j) \cap p(j+1)).$$

Since  $p' \cap p(j+1) \neq p(j) \cap p(j+1)$ ,

$$\dim(\pi(A_{p(j+1)}) \cap \pi(A_{p(j)}) \cap Z(t)) < k-1.$$

Also, since  $Z(t+1)$  is spanned by  $\pi(A_{p(j)}) \cup Z(t)$ ,

$$\dim(\pi(A_{p(j+1)}) \cap Z(t+1)) = k,$$

and hence

$$\pi(A_{p(j+1)}) \subset Z(t+1).$$

Therefore, (1) holds for  $p(j+1)$ . If  $q_{j+1} = k$ , interchange  $k$  and  $j+1$  in the argument above. This completes the proof of Part 1.

**PART 2.** As a consequence of Part 1, there is an  $H \in \mathcal{A}_n$  such that  $AH^k$  is diagonal. Hence we can assume that  $A$  is diagonal.

$$A = \text{diag}(a_p), \quad \text{for } p \in P(n, k).$$

Now select any two integers  $g$  and  $h$ , such that  $1 \leq g, h \leq n$ , and  $g \neq h$ . Let  $q$  and  $r$  be two elements of  $P(n, k-1)$ , neither of which contains  $g$  or  $h$ . Let

$$p = q \cup \{g\},$$

$$p' = q \cup \{h\},$$

$$\bar{p} = r \cup \{g\},$$

and

$$\bar{p}' = r \cup \{h\}.$$

We want to show that

$$a_p a_{\bar{p}} = a_{p'} a_{\bar{p}'}$$

As in Part 1, we construct two families of elements of  $P(n, k)$ .

$$p(0) = p,$$

$$p(j) = (p(j-1) - \{q_j\}) \cup \{r_j\},$$

for  $j = 1, 2, 3, \dots, k-1$  and

$$p'(0) = p',$$

$$p'(j) = (p'(j-1) - \{p'_j\}) \cup \{r_j\},$$

for  $j = 1, 2, 3, \dots, k-1$ . Here we regard the  $p(j)$  and  $p'(j)$  as so ordered that  $g$  or  $h$  is always the last element. It suffices to prove that

$$(2) \quad a_{p(j-1)} a_{p'(j)} = a_{p(j)} a_{p'(j-1)}$$

for  $j = 1, 2, 3, \dots, k-1$ . Let  $y = e_{p(j-1)} + e_{p(j)} + e_{p'(j)} + e_{p'(j-1)}$ . Then  $y \in \psi(k, n)$ . Therefore  $Ay \in \psi(k, n)$ . Thus  $Ay$  satisfies the Plucker identities, one of which may be written as (2), since only these four components of  $Ay$  are not zero. Now let

$$b(g, h) = a_p / a_{p'}.$$

Then  $b(g, h)$  is independent of  $q$ , and for any three integers  $g, h$ , and  $s$ ,  $1 \leq g, h, s \leq n$ ,

$$b(g, s) = b(g, h) b(h, s).$$

Therefore, for  $r \in P(n, k)$ , and  $r' = \{1, 2, 3, \dots, k\}$ ,

$$a_r = \prod_{i=1}^k b(p_i, i) a_{r'},$$

where  $\prod$  here indicates product. So if  $B \in \mathcal{A}_n$

$$B = \text{diag}(b(1, 1), b(2, 1), b(3, 1), \dots, b(n, 1)),$$

and

$$\lambda = \left( \prod_{i=1}^k b(1, i) \right) a_{r'},$$

then

$$A = \lambda B^k.$$

This completes the proof of Lemma 3.5.

6. **The orthogonal group.** In this section we let  $F$  be the field of real numbers. For  $m$  a positive integer, let  $\cdot$  denote the usual inner product of  $E_m$ , and  $|v|$  the usual norm. For  $A \in \mathcal{A}_m$ , let  $A^{(i)}$  denote the  $i$ th row vector of the matrix of  $A$ .

6.1. **LEMMA.** For  $m$  and  $A$  as above, if there exists a set  $T \subset E_m$  such that

1.  $e_i \in T$  for all integers  $i$ ,  $1 \leq i \leq m$ ,
2.  $A^{(i)} \in T$  for all integers  $i$ ,  $1 \leq i \leq m$ ,
3. for all  $v \in T$ ,  $Av \in T$ , and  $A^{-1}v \in T$ ,
4. for all  $v \in T$ ,  $|Av| = |v|$ ,

then  $A$  is orthonormal.

**Proof.** Since, for  $v \in T$ ,  $A^{-1}v \in T$ , we have that

$$|v| = |AA^{-1}(v)| = |A^{-1}(v)|.$$

Now let  $x_i = A^{-1}e_i$  for any integer  $i$ ,  $1 \leq i \leq m$ . Then  $|x_i| = 1$ , and  $Ax_i = e_i$ . Hence  $A^{(i)} \cdot x_i = 1$ , and thus  $|A^{(i)}| \geq 1$ . But

$$|AA^{(i)}|^2 = \sum_{j=1}^m (A^{(j)} \cdot A^{(i)})^2 = A^{(i)} \cdot A^{(i)}.$$

So

$$\sum_{j=1, j \neq i}^m (A^{(j)} \cdot A^{(i)})^2 = A^{(i)} \cdot A^{(i)}(1 - A^{(i)} \cdot A^{(i)}).$$

Hence  $|A^{(i)}| \leq 1$ . Thus, for any integers  $i$  and  $j$ ,  $1 \leq i, j \leq m$ ,  $i \neq j$ ,  $|A^{(i)}| = 1$ , and  $A^{(i)} \cdot A^{(j)} = 0$ . Hence  $A$  is orthonormal.

6.2. **THEOREM.** Let  $A \in G$  such that for all  $v \in \psi(k, n)$ ,  $|Av| = |v|$ . Then  $A$  is orthonormal, and there exist  $B \in \mathcal{A}_n$ ,  $B$  orthonormal, and  $C \in C_N$ ,  $C^2 = I$ , such that either  $A = CB^k$ , or  $A = CJB^k$ .

**Proof.** This follows immediately from the previous lemma.

#### REFERENCES

1. W. V. D. Hodge and D. Pedoe, *Methods of algebraic geometry*, The University Press, Cambridge, 1947-1952.
2. N. Bourbaki, *Eléments de mathématique*, II, Chapitre 3, *Algèbre multilinéaire*, Hermann, Paris, 1958.

SOUTHWESTERN AT MEMPHIS,  
MEMPHIS, TENNESSEE